

Contracting orbits in Outer space

Spencer Dowdall and Samuel J. Taylor

February 13, 2015

Abstract

We show that strongly contracting geodesics in Outer space project to parameterized quasi-geodesics in the free factor complex. This result provides a converse to a theorem of Bestvina–Feighn, and is used to give conditions for when a subgroup of $\text{Out}(\mathbb{F})$ has a quasi-isometric orbit map into the free factor complex.

1 Introduction

A geodesic $\gamma: \mathbf{I} \rightarrow X$ in a metric space X is strongly contracting if the closest point projection to γ contracts far away metric balls in X to sets of uniformly bounded diameter. Such geodesics exhibit hyperbolic-like behavior and are thus important to understanding the structure of the space. This note further develops the theory of strongly contracting geodesics in Outer space with the aim of understanding their behavior under the projection to the free factor complex. See §3 for precise definitions.

Such geometric questions in Outer space are often motivated by their analogs in Teichmüller space. In that setting, strongly contracting geodesics play an important role in our understanding of the geometry of Teichmüller space and the mapping class groups. These geodesics are characterized by the following result of Minsky describing both their structure in Teichmüller space and their behavior in the curve complex. (The equivalence of 1. and 2. in [Theorem 1.1](#) is the main result of [\[Min\]](#) while the equivalence of 1. and 3. follows easily from Theorem 4.3 of [\[Min\]](#).)

Theorem 1.1 (Minsky [\[Min\]](#)). *Let $\tau: \mathbf{I} \rightarrow \text{Teich}(S)$ be a Teichmüller geodesic. Then the following are equivalent:*

1. *There is an $\varepsilon \geq 0$ such that τ is entirely contained in the ε -thick part of $\text{Teich}(S)$.*
2. *There is a $D \geq 0$ such that τ is a D -strongly contracting geodesic in $\text{Teich}(S)$.*
3. *There is a $K \geq 1$ such that τ projects to a K -quasigeodesic in $\mathcal{C}(S)$, the curve complex of S .*

Moreover, the constants ε, D, K above depend only on each other and the topology of S .

Thus strongly contracting geodesics greatly illuminate the connection between Teichmüller space and the curve complex, as it is along these geodesics that the projection $\text{Teich}(S) \rightarrow \mathcal{C}(S)$ is quasi-isometric.

Our main result is a version of [Theorem 1.1](#) for the Outer space \mathcal{X} of a free group \mathbb{F} and its projection to the free factor complex \mathcal{F} of \mathbb{F} . This projection has already proven to be highly useful beginning with Bestvina and Feighn’s proof of hyperbolicity of the free factor complex [\[BF2\]](#) and continuing with, for example, [\[BR, Ham1, DT, Hor\]](#). In fact, in the course of proving hyperbolicity

of \mathcal{F} , Bestvina and Feighn show that folding path geodesics which made definite progress in \mathcal{F} are strongly contracting with respect to a specific projection tailored to foldings paths. Combining with Theorem 4.1 and Lemma 4.11 from [DT], this result of Bestvina and Feighn ([BF2, Corollary 7.3]) may be promoted to all geodesics:

Theorem 1.2. *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a geodesic whose projection to \mathcal{F} is a K -quasigeodesic. Then there exists $D \geq 0$ depending only on K (and the injectivity radius of the terminal endpoint of γ) such that γ is D -strongly contracting in \mathcal{X} .*

Here we prove a converse to **Theorem 1.2**. Together, these establish an analog of **Theorem 1.1** in the free group setting.

Theorem 1.3. *For each $D \geq 0$ there exists a constant $K \geq 1$ with the following property. If $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a nondegenerate D -strongly contracting geodesic, then $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a K -quasigeodesic.*

Recall that the Lipschitz metric on Outer space is not symmetric. Hence, a geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is not necessarily a (quasi) geodesic when traversed in the reverse direction. The condition that γ be **nondegenerate** in **Theorem 1.3** is, informally, that the backwards distance along γ meets a certain threshold depending only on D . See §3 for a precise definition and discussion.

Remark 1.4 (Thick geodesics in \mathcal{X}). Combining **Theorem 1.3** with **Theorem 1.2** gives the equivalence (2) \iff (3) of **Theorem 1.1** in the $\text{Out}(\mathbb{F})$ setting. We stress that the implication (1) \implies (2) of **Theorem 1.1** is in fact false in the Outer space setting. Indeed, it is well-known that there are thick geodesics in \mathcal{X} that nevertheless project to a bounded diameter subset of \mathcal{F} . By **Theorem 1.3** such geodesics cannot be strongly contracting.

Remark 1.5 (Hyperbolic isometries of \mathcal{F}). Combining **Theorem 1.3** with Algom-Kfir's result [AK] that axes of fully irreducible automorphisms in \mathcal{X} are strongly contracting gives an alternative proof of the fact that fully irreducible automorphisms act as loxodromic isometries on \mathcal{F} (i.e. they have positive translation length). This result was proven by Bestvina and Feighn in [BF2] using results in [BF1].

As an application to **Theorem 1.3**, we give conditions for when the orbit map from a finitely generated subgroup $\Gamma \leq \text{Out}(\mathbb{F})$ into \mathcal{F} is a quasi-isometric embedding. First, say that $\Gamma \leq \text{Out}(\mathbb{F})$ is **contracting in \mathcal{X}** if there exists $G \in \mathcal{X}$ and $D \geq 0$ so that for any two points in the orbit $\Gamma \cdot G$ there is a D -strongly contracting geodesic joining them.

Theorem 1.6. *Suppose that $\Gamma \leq \text{Out}(\mathbb{F})$ is finitely generated and that the orbit map $\Gamma \rightarrow \mathcal{X}$ is a quasi-isometric embedding. Then Γ is contracting in \mathcal{X} if and only if the orbit map $\Gamma \rightarrow \mathcal{F}$ to the free factor complex is a quasi-isometric embedding.*

We note that the “if” direction of **Theorem 1.6** appeared first in our earlier work [DT]. During the completion of the note, Hamenstädt and Hensel proved a related result [HH, Theorem 3]. Their theorem pertains to Morse geodesics in \mathcal{X} and relies on Hamenstädt's notion of lines of minima in \mathcal{X} , introduced in [Ham2]. We remark, however, that there is no a priori connection between strongly contracting and Morse geodesics in the asymmetric metric space \mathcal{X} without additional assumptions on the geodesic.

Acknowledgments. The first named author is partially supported by NSF grant DMS-1204814. The second named author is partially supported by NSF grant DMS-1204592. We also acknowledge support from NSF grants DMS 1107452, 1107263, 1107367 "RNMS: GEometric structures And Representation varieties" (the GEAR Network) and from the GATSBY seminar at Brown & Yale.

2 Background

We briefly recall the necessary background material on the metric structure of Outer space; see [FM, BF2, DT] for additional details.

Outer space. Let \mathbb{F} denote the free group of rank $r = \text{rk}(\mathbb{F}) \geq 3$. Let \mathcal{R} denote the r -petal rose with vertex $v \in \mathcal{R}$, and fix an isomorphism $\mathbb{F} \cong \pi_1(\mathcal{R}, v)$. For our purposes, a **graph** is a 1-dimensional CW complex, and a connected, simply connected graph is a **tree**. A **core graph** is a finite graph all of whose vertices have degree at least 2.

We now define Culler and Vogtmann's [CV] Outer space \mathcal{X} of marked metric graphs. A **marked graph** (G, g) is a core graph G together with a **marking** $g: \mathcal{R} \rightarrow G$, i.e. a homotopy equivalence. A **metric** on G is a function $\ell: E(G) \rightarrow \mathbb{R}_{>0}$ from the set of edges of G to the positive real numbers, which assigns a length $\ell(e)$ to each edge $e \in E(G)$. The sum $\sum_{e \in E(G)} \ell(e)$ is called the **volume** of G . With this setup, a **marked metric graph** is defined to be the triple (G, g, ℓ) ; two triples (G_1, g_1, ℓ_1) and (G_2, g_2, ℓ_2) are **equivalent** if there is a graph isometry $\phi: G_1 \rightarrow G_2$ that preserves the markings in the sense that $\phi \circ g_1$ is homotopic to g_2 . **Outer space**, denoted \mathcal{X} , is the set of equivalence classes of marked metric graphs of volume 1.

The marking $\mathcal{R} \rightarrow G$ for $G \in \mathcal{X}$ allows us to view any nontrivial conjugacy class α in \mathbb{F} as a homotopy class of loops in the core graph G . The unique immersed loop in this homotopy class is denoted by $\alpha|G$, which we view as an immersion of S^1 into G . The **length of α in $G \in \mathcal{X}$** , denoted $\ell(\alpha|G)$, is the sum of the lengths of the edges of G crossed by $\alpha|G$, counted with multiplicities. The **standard topology on \mathcal{X}** is the coarsest topology such that all of the length functions $\ell(\alpha|\cdot): \mathcal{X} \rightarrow \mathbb{R}_+$ are continuous [CV]. This topology agrees with other naturally defined topologies on \mathcal{X} , including the one induced by the Lipschitz metric defined below. See [CV, Pau, FM] for details.

A **difference of markings** from $G \in \mathcal{X}$ to $H \in \mathcal{X}$ is any map $\phi: G \rightarrow H$ that is homotopic to $h \circ g^{-1}$, where g and h are the markings on G and H , respectively. The **Lipschitz distance** from G to H is then defined to be

$$d_{\mathcal{X}}(G, H) := \inf\{\log(\text{Lip}(\phi)) \mid \phi \simeq h \circ g^{-1}\},$$

where $\text{Lip}(\phi)$ denotes the Lipschitz constant of the difference of markings ϕ . We note that while $d_{\mathcal{X}}$ is in general asymmetric (that is, $d_{\mathcal{X}}(G, H) \neq d_{\mathcal{X}}(H, G)$), it satisfies definiteness ($d_{\mathcal{X}}(G, H) = 0$ if and only if $G = H$) and the ordered triangle inequality ($d_{\mathcal{X}}(E, H) \leq d_{\mathcal{X}}(E, G) + d_{\mathcal{X}}(G, H)$) [FM]. We also have the following important result, originally due to Tad White, relating the Lipschitz distance to the ratio of lengths of conjugacy classes in the two graphs:

Proposition 2.1 (See Francaviglia–Martino [FM] or Algom-Kfir [AK]). *For every $G \in \mathcal{X}$ there exists a finite set \mathcal{C}_G of primitive conjugacy classes, called **candidates**, whose immersed representatives in G cross each edge at most twice and such that for any $H \in \mathcal{X}$*

$$d_{\mathcal{X}}(G, H) = \max_{\alpha \in \mathcal{C}_G} \log\left(\frac{\ell(\alpha|H)}{\ell(\alpha|G)}\right) = \sup_{\alpha \in \mathbb{F}} \log\left(\frac{\ell(\alpha|H)}{\ell(\alpha|G)}\right).$$

Note that because each candidate $\alpha \in \mathcal{C}_G$ crosses each edge of G no more than twice, $\ell(\alpha|G) \leq 2$.

Finally, a **geodesic in \mathcal{X}** is by definition a *directed* geodesic, that is, a path $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ such that $d_{\mathcal{X}}(\gamma(s), \gamma(t)) = t - s$ for all $s < t$. Throughout, \mathbf{I} will always denote a closed interval $\mathbf{I} \subset \mathbb{R}$, and we write $\mathbf{I}_{\pm} \in \mathbb{R} \cup \{\pm\infty\}$ for the (possibly infinite) endpoints of the interval \mathbf{I} .

Asymmetry and the thick part of Outer space. For $\varepsilon > 0$, we define the ε -**thick part** of \mathcal{X} to be the subset

$$\mathcal{X}_\varepsilon := \{G \in \mathcal{X} : \ell(\alpha|G) \geq \varepsilon \text{ for every nontrivial conjugacy class } \alpha \text{ in } \mathbb{F}\}.$$

It is also sometimes convenient to consider the symmetrization of the Lipschitz metric:

$$d_{\mathcal{X}}^{\text{sym}}(G, H) := d_{\mathcal{X}}(G, H) + d_{\mathcal{X}}(H, G)$$

which is an actual metric on \mathcal{X} and induces the standard topology [FM]. Because the Lipschitz metric $d_{\mathcal{X}}$ is not symmetric, care must be taken when discussing distances in \mathcal{X} . This asymmetry, however, is somewhat controlled in the thick part \mathcal{X}_ε :

Lemma 2.2 (Handel–Mosher [HM], Algom-Kfir–Bestvina [AKB]). *For any $\varepsilon > 0$, there exists $M_\varepsilon \geq 1$ so that for all $G, H \in \mathcal{X}_\varepsilon$ we have*

$$d_{\mathcal{X}}(H, G) \leq d_{\mathcal{X}}^{\text{sym}}(H, G) = d_{\mathcal{X}}^{\text{sym}}(G, H) \leq M_\varepsilon \cdot d_{\mathcal{X}}(G, H).$$

For $G, H \in \mathcal{X}$, we also use the notation $\text{diam}_{\mathcal{X}}(G, H)$ to denote $\text{diam}_{\mathcal{X}}\{G, H\} = \max\{d_{\mathcal{X}}(G, H), d_{\mathcal{X}}(H, G)\}$. Observe that $\text{diam}_{\mathcal{X}}(G, H) \leq d_{\mathcal{X}}^{\text{sym}}(G, H) \leq 2\text{diam}_{\mathcal{X}}(G, H)$.

The factor complex. This main purpose of this note is to show that strongly contracting geodesics in \mathcal{X} project to parameterized quasigeodesics in the free factor complex, which is defined as follows: The **factor complex** \mathcal{F} associated to the free group \mathbb{F} is the simplicial complex whose vertices are conjugacy classes of proper, nontrivial free factors of \mathbb{F} . Vertices $[A_0], \dots, [A_k]$ span a k -simplex if, after reordering, we have the proper inclusions $A_0 < \dots < A_k$. This simplicial complex was first introduced by Hatcher and Vogtmann in [HV]. Since we are interested in the coarse geometry of \mathcal{F} , we will only consider its 1-skeleton equipped with the path metric induced by giving each edge length 1. The following theorem of Bestvina and Feighn is fundamental to the geometric study of $\text{Out}(\mathbb{F})$:

Theorem 2.3 (Bestvina–Feighn [BF2]). *The factor complex \mathcal{F} is Gromov hyperbolic.*

The primitive loop complex. For our proof of [Theorem 1.3](#), it is more natural to work with a different complex that is nevertheless quasi-isometric to \mathcal{F} . Recall that an element $\alpha \in \mathbb{F}$ is **primitive** if it is part of some free basis of \mathbb{F} . Thus α is primitive if and only if α generates a cyclic free factor of \mathbb{F} . We use the terminology **primitive loop** to mean a conjugacy class of \mathbb{F} consisting of primitive elements. The **primitive loop complex** \mathcal{PL} is then defined to be the simplicial graph whose vertices are primitive loops and where two vertices are joined by an edge in \mathcal{PL} if their respective conjugacy classes have representatives that are jointly part of a free basis of \mathbb{F} . It is straightforward to show that the natural inclusion map $\mathcal{PL}^0 \rightarrow \mathcal{F}^0$ (each primitive conjugacy class is itself a free factor) is 2-biLipschitz. Since the image is 1-dense, this map is in fact a 2-quasi-isometry.

Relating Outer space to the primitive loop graph, we define the **projection** $\pi_{\mathcal{PL}} : \mathcal{X} \rightarrow \mathcal{PL}$ in the following way: For $G \in \mathcal{X}$, set

$$\pi_{\mathcal{PL}}(G) := \{\alpha \in \mathcal{PL} : \ell(\alpha|G) \leq 2\}.$$

This is, of course, closely related to the projection $\pi_{\mathcal{F}} : \mathcal{X} \rightarrow \mathcal{F}$ defined by Bestvina and Feighn in [BF2] sending $G \in \mathcal{X}$ to the collection of free factors corresponding to proper core subgraphs of G . They prove that $\text{diam}_{\mathcal{F}}(\pi_{\mathcal{F}}(G)) \leq 4$ [BF2, Lemma 3.1] and that $\text{diam}_{\mathcal{F}}(\alpha \cup \pi_{\mathcal{F}}(G)) \leq 6\ell(\alpha|G) + 13$

[BF2, Lemma 3.3] for every $G \in \mathcal{X}$ and every primitive conjugacy class α . These estimates imply that $\pi_{\mathcal{P}\mathcal{L}}$ and $\pi_{\mathcal{F}}$ coarsely agree under the 2–quasi-isometry $\mathcal{P}\mathcal{L} \rightarrow \mathcal{F}$ defined above. Combining with the fact that $\pi_{\mathcal{F}}$ is coarsely Lipschitz [BF2, Corollary 3.5] this moreover gives the existence of a constant $L \geq 1$ such that

$$d_{\mathcal{P}\mathcal{L}}(G, H) := \text{diam}_{\mathcal{P}\mathcal{L}}(\pi_{\mathcal{P}\mathcal{L}}(G) \cup \pi_{\mathcal{P}\mathcal{L}}(H)) \leq L d_{\mathcal{X}}(G, H) + L$$

for all $G, H \in \mathcal{X}$. That is, the projection $\pi_{\mathcal{P}\mathcal{L}} : \mathcal{X} \rightarrow \mathcal{P}\mathcal{L}$ is coarsely L –Lipschitz. Its easily computed that $L \leq 260$, but we prefer to work with the symbol L for clarity.

3 Strongly contracting geodesics

Suppose that $\gamma : \mathbf{I} \rightarrow \mathcal{X}$ is a (directed) geodesic. Then for any point $H \in \mathcal{X}$ we write $d_{\mathcal{X}}(H, \gamma) = \inf\{t \in \mathbf{I} \mid d_{\mathcal{X}}(H, \gamma(t))\}$ for the infimal distance from H to γ . The **closest point projection of H to γ** is then defined to be the set

$$\pi_{\gamma}(H) := \{\gamma(t) \mid t \in \mathbf{I} \text{ such that } d_{\mathcal{X}}(H, \gamma(t)) = d_{\mathcal{X}}(H, \gamma)\} \subset \mathcal{X}.$$

Note that $\pi_{\gamma}(H)$ could in principle have infinite diameter: due to the asymmetry of $d_{\mathcal{X}}$, the directed triangle inequality does not in general allow one to bound $d_{\mathcal{X}}(\gamma(s), \gamma(t))$ for times $s < t$ with $\gamma(s), \gamma(t) \in \pi_{\gamma}(H)$. On the other extreme, it is conceivable that the above infimum need not be realized (since $d_{\mathcal{X}}(H, \gamma(s))$ could remain bounded as $s \rightarrow -\infty$), in which case $\pi_{\gamma}(H) = \emptyset$ by definition.

Definition 3.1 (Strongly contracting). *A geodesic $\gamma : \mathbf{I} \rightarrow \mathcal{X}$ is D –strongly contracting if for all points $H, H' \in \mathcal{X}$ satisfying $d_{\mathcal{X}}(H, H') \leq d_{\mathcal{X}}(H, \gamma)$ we have*

- $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H')) \leq D$, and
- $\pi_{\gamma}(H) = \emptyset$ if and only if $\pi_{\gamma}(H') = \emptyset$.

We say that γ is strongly contracting if it is D –strongly contracting for some $D \geq 0$.

We remark that the second condition is a natural extension of the first: $\pi_{\gamma}(H) = \emptyset$ only if there is a sequence $s_i \in \mathbf{I}$ tending to $\mathbf{I}_- = -\infty$ with $d_{\mathcal{X}}(H, \gamma(s_i))$ limiting to $d_{\mathcal{X}}(H, \gamma)$. In this case, one should morally view $\pi_{\gamma}(H)$ as being “ $\gamma(-\infty)$ ”; hence $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H'))$ is considered to be infinite unless $\pi_{\gamma}(H') = \emptyset$ as well.

While $\pi_{\gamma}(H)$ may in principle be empty, our first lemma shows that the closest point projection $\pi_{\gamma}(H)$ always exists when γ is strongly contracting:

Lemma 3.2. *If $\gamma : \mathbf{I} \rightarrow \mathcal{X}$ is D –strongly contracting, then $\pi_{\gamma}(H)$ is nonempty for all $H \in \mathcal{X}$.*

Proof. Let us first show that $\pi_{\gamma}(H) \neq \emptyset$ for all H in an open neighborhood of $\gamma(\mathbf{I})$. Let $t \in \mathbf{I}$ be arbitrary and let $U \subset \mathbf{I}$ be an open neighborhood of t whose closure \bar{U} is a compact, proper subinterval of \mathbf{I} . Then there exists $C > 0$ such that $|s - t| \geq C$ and consequently $d_{\mathcal{X}}^{\text{sym}}(\gamma(t), \gamma(s)) \geq C$ for all $s \in \mathbf{I} \setminus U$. In particular, we have $\gamma(s) \neq \gamma(t)$ and thus $d_{\mathcal{X}}(\gamma(t), \gamma(s)) > 0$ for all $s \in \mathbf{I} \setminus U$. Moreover, it is easily shown that the infimum

$$\varepsilon_t = \inf\{d_{\mathcal{X}}(\gamma(t), \gamma(s)) : s \in \mathbf{I} \setminus U\}$$

is in fact positive.

Now consider any point H in the open neighborhood $V_t = \{y \in \mathcal{X} : d_{\mathcal{X}}^{\text{sym}}(y, \gamma(t)) < \varepsilon_t/3\}$ of $\gamma(t)$. If $d_{\mathcal{X}}(H, \gamma(s)) \leq \varepsilon_t/2$ for some $s \in \mathbf{I} \setminus U$, the triangle inequality would give

$$d_{\mathcal{X}}(\gamma(t), \gamma(s)) \leq d_{\mathcal{X}}^{\text{sym}}(\gamma(t), H) + d_{\mathcal{X}}(H, \gamma(s)) < \varepsilon_t/3 + \varepsilon_t/2 < \varepsilon_t,$$

which is impossible by definition of ε_t . Hence $d_{\mathcal{X}}(H, \gamma(s)) \geq \varepsilon_t/2$ for all $s \in \mathbf{I} \setminus U$. Since $d_{\mathcal{X}}(H, \gamma) \leq d_{\mathcal{X}}(H, \gamma(t)) < \varepsilon_t/3$, it follows that

$$d_{\mathcal{X}}(H, \gamma) = \inf\{i \in \mathbf{I} : d_{\mathcal{X}}(H, \gamma(i))\} = \inf\{s \in \bar{U} : d_{\mathcal{X}}(H, \gamma(s))\}.$$

The above infimum is necessarily realized by compactness; thus we conclude $d_{\mathcal{X}}(H, \gamma) = d_{\mathcal{X}}(H, \gamma(s))$ for some $s \in \bar{U}$. This proves that $\pi_{\gamma}(H) \neq \emptyset$ for all points H in the open neighborhood $V = \cup_{t \in \mathbf{I}} V_t$ of $\gamma(\mathbf{I})$. Note that we have not yet used the assumption that γ is strongly contracting.

Let us now employ strong contraction to complete the proof of the lemma. Let $H \in \mathcal{X}$ be arbitrary; we may assume $H \notin \gamma(\mathbf{I})$ for otherwise the claim is obvious. Choose any path $\mu : [a, b] \rightarrow \mathcal{X}$ from $\mu(a) = H$ to some point $\mu(b) \in \gamma(\mathbf{I})$. By restricting to a smaller interval if necessary, we may additionally assume that $\mu(s) \notin \gamma(\mathbf{I})$ for all $a \leq s < b$. Since $\pi_{\gamma}(G)$ is nonempty for all G in a neighborhood of $\gamma(\mathbf{I})$, there exists some $c \in (a, b)$ such that $\pi_{\gamma}(\mu(c))$ is nonempty.

Now, for each $t \in [a, c]$ we have $\mu(t) \notin \gamma(\mathbf{I})$. In fact we claim that $d_{\mathcal{X}}(\mu(t), \gamma) > 0$: otherwise, as above, there would exist a sequence $s_i \in \mathbf{I}$ with $d_{\mathcal{X}}(\mu(t), \gamma(s_i)) \rightarrow 0$ and consequently $\gamma(s_i) \rightarrow \mu(t)$, contradicting the fact that $\gamma(\mathbf{I})$ is closed. Thus for each $t \in [a, c]$ we may find an open neighborhood $W_t \subset \mathcal{X}$ of $\mu(t)$ such that $d_{\mathcal{X}}(\mu(t), G') \leq d_{\mathcal{X}}(\mu(t), \gamma)$ for all $G' \in W_t$. By compactness, there is a finite subcollection W_1, \dots, W_k of these open sets that cover $\mu([a, c])$. For each i the strongly contracting condition now implies that either $\pi_{\gamma}(G') = \emptyset$ for all $G' \in W_i$, or else $\pi_{\gamma}(G') \neq \emptyset$ for all $G' \in W_i$. Since these W_1, \dots, W_k cover the connected set $\mu([a, c])$ and at least one W_i falls into the latter category (namely, the set W_i containing $\mu(c)$), the contingency “ $\pi_{\gamma}(G') \neq \emptyset$ for all $G' \in W_i$ ” must in fact hold for every i . In particular, we see that $\pi_{\gamma}(H) = \pi_{\gamma}(\mu(a))$ is nonempty, as claimed. \square

We will also need the following basic observation showing that the projection of a connected set to a D -strongly contracting geodesic γ is effectively “ D -dense” in $\gamma(\mathbf{I})$:

Lemma 3.3. *Let $\gamma : \mathbf{I} \rightarrow \mathcal{X}$ be D -strongly contracting and let $[a, b] \subset \mathbf{I}$ be any subinterval with $|a - b| \geq D$. If $A \subset \mathcal{X}$ is connected and $\pi_{\gamma}(A)$ misses $\gamma([a, b])$ (that is $\pi_{\gamma}(A) \cap \gamma([a, b]) = \emptyset$), then $\pi_{\gamma}(A)$ is either entirely contained in $\gamma(\mathbf{I} \cap (-\infty, a))$ or entirely contained in $\gamma(\mathbf{I} \cap (b, \infty))$.*

Proof. Let us first establish the following

Claim. *Each $H \in \mathcal{X}$ admits a neighborhood $U \subset \mathcal{X}$ with $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H')) \leq D$ for all $H' \in U$.*

To prove the claim, first suppose $H \notin \gamma(\mathbf{I})$ so that, as above, we have $\delta := d_{\mathcal{X}}(H, \gamma) > 0$ (for otherwise there is a sequence in $\gamma(\mathbf{I})$ converging to H). Taking the open neighborhood to be $U = \{y \in \mathcal{X} : d_{\mathcal{X}}(H, y) < \delta\}$, the strongly contracting condition then ensures $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H), \pi_{\gamma}(H')) \leq D$ for all $H' \in U$. Next suppose $H \in \gamma(\mathbf{I})$ so that $d_{\mathcal{X}}(H, \gamma) = 0$ and $\pi_{\gamma}(H) = \{H\}$. Choosing $\varepsilon > 0$ so that $H \in \mathcal{X}_{2\varepsilon}$, we may then choose $\delta > 0$ sufficiently small so that $\delta M_{\varepsilon} < D/2$ and the entire neighborhood $U = \{y \in \mathcal{X} : d_{\mathcal{X}}^{\text{sym}}(y, H) < \delta\}$ is contained in $\mathcal{X}_{\varepsilon}$ (where M_{ε} is the symmetrization constant from [Lemma 2.2](#)). For any $H' \in U$ we then have $d_{\mathcal{X}}(H', \gamma) \leq d_{\mathcal{X}}(H', H) < \delta$. Thus any $G \in \pi_{\gamma}(H')$ satisfies $d_{\mathcal{X}}^{\text{sym}}(G, H') \leq M_{\varepsilon} d_{\mathcal{X}}(H', G) < M_{\varepsilon} \delta < D/2$. Note that we also have $d_{\mathcal{X}}^{\text{sym}}(H, H') < \delta < D/2$. Since $\pi_{\gamma}(H) = \{H\}$, the triangle inequality therefore shows the desired inequality $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H')) \leq D$. Since this holds for each $H' \in U$, the claim follows.

We now prove the lemma. Let $[a, b] \subset \mathbf{I}$ and $A \subset \mathcal{X}$ be as in the statement of the lemma, so that $\pi_{\gamma}(A)$ is disjoint from $\gamma([a, b])$. Since $\pi_{\gamma}(H)$ is always nonempty (by [Lemma 3.2](#)) and satisfies

$\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H)) \leq D$, the hypothesis on $\pi_{\gamma}(A) \cap \gamma([a, b])$ implies that each $H \in A$ lies in exactly one of two the sets

$$A_- = \{H \in A : \pi_{\gamma}(H) \subset \gamma(\mathbf{I} \cap (-\infty, a))\} \quad \text{or} \quad A_+ = \{H \in A : \pi_{\gamma}(H) \subset \gamma(\mathbf{I} \cap (b, \infty))\}.$$

Thus $A = A_- \cup A_+$ gives a partition of A . Moreover, the claim proves that A_- and A_+ are both open. The connectedness of A therefore implies that either A_- or A_+ is empty, which is exactly the conclusion of the lemma. \square

We say that a D -strongly contracting geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is **nondegenerate** if there exists times $s < t$ in \mathbf{I} such that $d_{\mathcal{X}}(\gamma(t), \gamma(s)) \geq 18DL$. The following lemma shows that this mild symmetry condition automatically holds in most natural situations. The proof is given in §7 and will follow easily from the tools developed in §§4–6.

Lemma 3.4. *Suppose that $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a D -strongly contracting geodesic. Then either of the following conditions imply that γ is nondegenerate:*

- $|\mathbf{I}| \geq A$ for some constant A depending only on D and the injectivity radius of $\gamma(\mathbf{I}_-)$.
- \mathbf{I} is an infinite length interval

4 Length minimizers

To control the nearest point projection of a graph H to a geodesic γ , we must understand where the lengths of conjugacy classes in \mathbb{F} are minimized along γ . To this end, we introduce the following terminology: Firstly, given a directed geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ and a nontrivial conjugacy class $\alpha \in \mathbb{F}$, we typically write

$$m_{\alpha} = m_{\alpha}^{\gamma} := \inf_{t \in \mathbf{I}} \ell(\alpha | \gamma(t))$$

for the infimal length that the conjugacy class attains along γ . We then regard the set

$$\rho_{\gamma}(\alpha) = \{t \in \mathbf{I} \mid \ell(\alpha | \gamma(t)) = m_{\alpha}\}$$

as the **projection of α to γ** . Since it is possible to have $\rho_{\gamma}(\alpha) = \emptyset$ in the case that \mathbf{I} is not compact, we also define a parameterwise-projection

$$\hat{\rho}'_{\gamma}(\alpha) = \{t \mid \exists s_i \in \mathbf{I} \text{ s.t. } s_i \rightarrow t \text{ and } \ell(\alpha | \gamma(s_i)) \rightarrow m_{\alpha}\} \subset [-\infty, +\infty].$$

For technical reasons, it is convenient to instead work with the following variant:

$$\hat{\rho}_{\gamma}(\alpha) := \begin{cases} \hat{\rho}'_{\gamma}(\alpha) \cap \mathbb{R}, & \rho_{\gamma}(\alpha) \neq \emptyset \\ \hat{\rho}'_{\gamma}(\alpha), & \text{else.} \end{cases}$$

Thus $\hat{\rho}_{\gamma}(\alpha)$ is never empty and is exactly the set of parameters $t \in \mathbf{I}$ realizing m_{α} when $\rho_{\gamma}(\alpha)$ is nonempty. Note also that $\gamma(\hat{\rho}_{\gamma}(\alpha) \cap \mathbb{R}) = \rho_{\gamma}(\alpha)$ in all cases.

As indicated above, we think of ρ_{γ} as a projection from the set of conjugacy classes in \mathbb{F} onto γ . The next lemma shows that for strongly contracting geodesics, ρ_{γ} is compatible with closest-point projection π_{γ} in the sense that graphs $H \in \mathcal{X}$ and embedded loops in H often coarsely project to the same spot.

Lemma 4.1 (Projections agree). *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a D -strongly contracting geodesic. Suppose that $H \in \mathcal{X}$ is such that $d_{\mathcal{X}}(H, \gamma) \geq \log(3)$. Then for every conjugacy class α corresponding to an embedded loop in H with $\ell(\alpha|H) \leq 2/3$, we have that $\rho_{\gamma}(\alpha) \neq \emptyset$ and that*

$$\text{diam}_{\mathcal{X}}(\rho_{\gamma}(\alpha) \cup \pi_{\gamma}(H)) \leq D.$$

Moreover, there exists a primitive conjugacy class α satisfying these conditions.

Proof. **Lemma 3.2** ensures the existence of a time $t \in \mathbf{I}$ so that $d_{\mathcal{X}}(H, \gamma) = d_{\mathcal{X}}(H, \gamma(t))$. Write $G = \gamma(t)$. Let α be a conjugacy class corresponding to an embedded loop in H with $\ell(\alpha|H) \leq 2/3$. Notice that every metric graph in \mathcal{X} indeed contains an embedded loop of length at most $2/3$.

To prove that α satisfies the conclusion of the lemma, for $0 < \sigma \leq 1$, let H_{σ} denote the metric graph obtained from H by scaling the edges comprising $\alpha|H \subset H$ by σ and scaling all other edges by $\frac{1-\sigma\ell(\alpha|H)}{1-\ell(\alpha|H)}$ (so as to maintain $\text{vol}(H_{\sigma}) = 1$). It follows that

$$d_{\mathcal{X}}(H, H_{\sigma}) \leq \log\left(\frac{1-\sigma\ell(\alpha|H)}{1-\ell(\alpha|H)}\right) \leq \log\left(\frac{1}{1-\ell(\alpha|H)}\right) \leq \log(3).$$

Since γ is D -strongly contracting and $d_{\mathcal{X}}(H, \gamma) \geq \log(3)$ by hypothesis, we may conclude that $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H_{\sigma})) \leq D$ for all $0 < \sigma \leq 1$.

First suppose that $\rho_{\gamma}(\alpha) \neq \emptyset$. Letting $B \in \rho_{\gamma}(\alpha)$ be arbitrary, we then have $\ell(\alpha|B) = m_{\alpha}$. Let \mathcal{C} be the set of candidates of H ; this is also the set of candidates for each H_{σ} . Since α is embedded in H , it is the only candidate whose edge lengths all tend to zero as $\sigma \rightarrow 0$. Thus for every candidate $z \neq \alpha \in \mathcal{C}$, there is a positive lower bound on $\ell(z|H_{\sigma})$ as $\sigma \rightarrow 0$. On the other hand, $\ell(\alpha|H_{\sigma})$ clearly tends to zero as $\sigma \rightarrow 0$. For $\sigma > 0$ sufficiently small, **Proposition 2.1** therefore gives

$$\log\left(\frac{m_{\alpha}}{\ell(\alpha|H_{\sigma})}\right) = \log\left(\frac{\ell(\alpha|B)}{\ell(\alpha|H_{\sigma})}\right) = \log\left(\max_{z \in \mathcal{C}} \frac{\ell(z|B)}{\ell(z|H_{\sigma})}\right) = d_{\mathcal{X}}(H_{\sigma}, B).$$

The fact that m_{α} is the minimal length achieved by α along γ moreover implies that $d_{\mathcal{X}}(H_{\sigma}, \gamma) \geq \log(m_{\alpha}/\ell(\alpha|H_{\sigma}))$. Therefore $d_{\mathcal{X}}(H_{\sigma}, B) = d_{\mathcal{X}}(H_{\sigma}, \gamma)$, and so we may conclude $B \in \pi_{\gamma}(H_{\sigma})$. This shows that $\rho_{\gamma}(\alpha) \subset \pi_{\gamma}(H_{\sigma})$ and therefore that $\text{diam}_{\mathcal{X}}(\rho_{\gamma}(\alpha) \cup \pi_{\gamma}(H)) \leq D$ in the case that $\rho_{\gamma}(\alpha)$ is nonempty.

It remains to rule out the possibility that $\rho_{\gamma}(\alpha)$ is empty. Since in this case the infimal length m_{α} is *only* achieved by sequences of times tending to $\pm\infty$, we may choose $\varepsilon > 0$ sufficiently small so that for any $s \in \mathbf{I}$ we have the implication

$$\ell(\alpha|\gamma(s)) < m_{\alpha} + \varepsilon \implies |s - t| > 2D.$$

Let us choose such a time $s_0 \in \mathbf{I}$ with $\ell(\alpha|\gamma(s_0)) < m_{\alpha} + \varepsilon$. As above, by taking $\sigma > 0$ sufficiently small we may be assured that

$$d_{\mathcal{X}}(H_{\sigma}, \gamma(s_0)) = \log\left(\max_{z \in \mathcal{C}} \frac{\ell(z|\gamma(s_0))}{\ell(z|H_{\sigma})}\right) = \log\left(\frac{\ell(\alpha|\gamma(s_0))}{\ell(\alpha|H_{\sigma})}\right).$$

Since $\pi_{\gamma}(H_{\sigma})$ is nonempty by **Lemma 3.2**, there exists a time $s \in \mathbf{I}$ for which $\gamma(s) \in \pi_{\gamma}(H_{\sigma})$. Since this is a *closest* point on γ from H_{σ} , we necessarily have $d_{\mathcal{X}}(H_{\sigma}, \gamma(s)) \leq d_{\mathcal{X}}(H_{\sigma}, \gamma(s_0))$ which in turn requires $\ell(\alpha|\gamma(s)) \leq \ell(\alpha|\gamma(s_0)) < m_{\alpha} + \varepsilon$. By the choice of ε , this implies $|s - t| > 2D$ and consequently $\text{diam}_{\mathcal{X}}(\gamma(s), \gamma(t)) > 2D$. However, since $\gamma(s) \in \pi_{\gamma}(H_{\sigma})$ and $\gamma(t) \in \pi_{\gamma}(H)$, this contradicts the fact that $\text{diam}_{\mathcal{X}}(\pi_{\gamma}(H) \cup \pi_{\gamma}(H_{\sigma})) \leq D$ for all $0 < \sigma \leq 1$. Therefore $\rho_{\gamma}(\alpha)$ cannot be empty, and the lemma holds. \square

A priori, it could be that $\rho_\gamma(\alpha)$ is empty for every nontrivial conjugacy class in \mathbb{F} and, in keeping with [Lemma 4.1](#), that all points of \mathcal{X} lies within $\log(3)$ of $\gamma(\mathbf{I})$. Our next lemma rules out such pathological behavior.

Lemma 4.2 (Some projection exists). *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a strongly contracting geodesic. Then there exists a primitive conjugacy class $\alpha \in \mathcal{P}\mathcal{L}^0$ such that $\rho_\gamma(\alpha)$ is nonempty and $m_\alpha > 0$.*

Proof. Suppose on the contrary that $\rho_\gamma(\alpha)$ is empty for all $\alpha \in \mathcal{P}\mathcal{L}^0$. Note that this requires the interval $\mathbf{I} \subset \mathbb{R}$ to have infinite length. We first claim that $m_\alpha = 0$ for all $\alpha \in \mathcal{P}\mathcal{L}^0$. Otherwise we may find some $\alpha \in \mathcal{P}\mathcal{L}^0$ with m_α positive, and we let $R \in \mathcal{X}$ be any rose with one petal labeled by α with $\ell(\alpha|R) = m_\alpha/3$. Then by definition of m_α , for every $t \in \mathbf{I}$ we have

$$d_{\mathcal{X}}(R, \gamma(t)) \geq \log\left(\frac{\ell(\alpha|\gamma(t))}{\ell(\alpha|R)}\right) \geq \log\left(\frac{m_\alpha}{m_\alpha/3}\right) = \log(3).$$

Since α corresponds to an embedded loop in R by construction, [Lemma 4.1](#) then gives $\rho_\gamma(\alpha) \neq \emptyset$, contradicting our assumption. Hence m_α is indeed zero for all $\alpha \in \mathcal{P}\mathcal{L}^0$.

Next, for each $\alpha \in \mathcal{P}\mathcal{L}^0$ we claim that $\{-\infty\} \in \hat{\rho}_\gamma(\alpha)$, i.e., that there exists a sequence $s_i \in \mathbf{I}$ with $s_i \rightarrow -\infty$ such that $\ell(\alpha|\gamma(s_i)) \rightarrow 0$. To see this, fix a rose R with one petal labeled by α , and for $0 < \sigma < 1$ let R_σ denote R equipped with the metric for which $\ell(\alpha|R) = \sigma$ and all other petals have length $\frac{1-\sigma}{r-1}$ (so that $\text{vol}(R_\sigma) = 1$). Fix also a parameter $t_0 \in \mathbf{I}$. Since $m_\alpha = 0$, for any given $\varepsilon_0 > 0$ we may choose $\varepsilon < \varepsilon_0$ sufficiently small such that the implication

$$\ell(\alpha|\gamma(s)) < \varepsilon \implies |s - t_0| > 2D$$

holds for every $s \in \mathbf{I}$. Moreover, since $m_\alpha = 0$ we may choose a point $H \in \gamma(\mathbf{I})$ with $\ell(\alpha|H) < \varepsilon$. We may also choose $\sigma_0 < 1$ sufficiently small so that for all $0 < \sigma \leq \sigma_0$ the candidate α of R_σ realizes the distance to H , that is

$$d_{\mathcal{X}}(R_\sigma, H) = \log\left(\frac{\ell(\alpha|H)}{\ell(\alpha|R_\sigma)}\right).$$

It follows that each point $G = \gamma(s) \in \pi_\gamma(R_\sigma)$ (with $0 < \sigma \leq \sigma_0$) satisfies

$$\log\left(\frac{\ell(\alpha|G)}{\ell(\alpha|R_\sigma)}\right) \leq d_{\mathcal{X}}(R_\sigma, G) \leq d_{\mathcal{X}}(R_\sigma, H) = \log\left(\frac{\ell(\alpha|H)}{\ell(\alpha|R_\sigma)}\right)$$

and consequently $\ell(\alpha|G) < \ell(\alpha|H) < \varepsilon$ and thus $|s - t_0| > 2D$. Since $\varepsilon > 0$ was chosen arbitrarily, this shows the maximal length of α on the set $\pi_\gamma(R_\sigma)$ tends to 0 as $\sigma \rightarrow 0$ and that the sets $\pi_\gamma(R_\sigma)$ leave every compact subset of $\gamma(\mathbf{I})$ as $\sigma \rightarrow 0$.

We additionally see that $\pi_\gamma(R_\sigma)$ is disjoint from the interval $\gamma(\mathbf{I} \cap [t_0 - 2D, t_0 + 2D])$ for all $0 < \sigma \leq \sigma_0$. Since $\mathbf{I} \cap [t_0 - 2D, t_0 + 2D]$ has length at least $2D$ (recall that \mathbf{I} must have infinite length) and the set $\mathcal{R} = \{R_\sigma : 0 < \sigma \leq \sigma_0\}$ is connected, [Lemma 3.3](#) shows that $\pi_\gamma(\mathcal{R})$ is either contained entirely within $\gamma(\mathbf{I} \cap (-\infty, t_0 + 2D])$ or entirely within $\gamma(\mathbf{I} \cap [t_0 + 2D, \infty))$. We claim that the former possibility must hold. Indeed, suppose instead that $\pi_\gamma(\mathcal{R})$ is contained in $\gamma(\mathbf{I} \cap [t_0 + 2D, \infty))$. For each $0 < \sigma \leq \sigma_0$, choose a parameter $s_\sigma \in \mathbf{I}$ such that $\gamma(s_\sigma) \in \pi_\gamma(R_\sigma)$; note that we necessarily have $s_\sigma > t_0 + 2D$ by our assumption. Since the sets $\pi_\gamma(R_\sigma)$ exit all compact subset of $\gamma(\mathbf{I})$, it follows that $s_\sigma \rightarrow \infty$ as $\sigma \rightarrow 0$. Let L_0 denote the distance $d_{\mathcal{X}}(\gamma(t_0), R_{\sigma_0})$. A comparison of lengths of candidates easily shows that the distance $d_{\mathcal{X}}(R_{\sigma_0}, R_\sigma)$ is bounded by $\log(3)$ for all $0 < \sigma \leq \sigma_0$. Furthermore,

we evidently have $d_{\mathcal{X}}(R_{\sigma}, \gamma) < \log(3)$, for otherwise [Lemma 4.1](#) would give $\rho_{\gamma}(\alpha) \neq \emptyset$ which is not the case. The triangle inequality therefore gives

$$s_{\sigma} - t_0 = d_{\mathcal{X}}(\gamma(t_0), \gamma(s_{\sigma})) \leq L_0 + 2 \log(3)$$

for all $0 < \sigma \leq \sigma_0$, contradicting the above observation that s_{σ} tends to ∞ . Thus the former possibility indeed holds and we may conclude that $\pi_{\gamma}(\mathcal{R})$ is contained in $\gamma(\mathbf{I} \cap (-\infty, t_0 - 2D])$. Therefore, choosing $s_{\sigma} \in \mathbf{I}$ so that $\gamma(s_{\sigma}) \in \pi_{\gamma}(R_{\sigma})$ as above, the preceding paragraph shows that $s_{\sigma} \rightarrow -\infty$ and $\ell(\alpha|s_{\sigma}) \rightarrow 0$ as $\sigma \rightarrow 0$. Thus $\{-\infty\} \in \hat{\rho}_{\gamma}(\alpha)$ as claimed.

We have so far shown that $m_{\alpha} = 0$ and $\{-\infty\} \in \hat{\rho}_{\gamma}(\alpha)$ for all $\alpha \in \mathcal{P}\mathcal{L}^0$. We will now use these facts to derive a contradiction and so prove the lemma. Fix $\alpha \in \mathcal{P}\mathcal{L}^0$ and for $0 < \sigma < 1$ let R_{σ} be the rose as above so that the projections $\pi_{\gamma}(R_{\sigma})$ exit the left end of $\gamma(\mathbf{I})$ as $\sigma \rightarrow 0$. Choose a primitive loop $\beta \in \mathcal{P}\mathcal{L}^0$ with $d_{\mathcal{F}}(\alpha, \beta) > 30$. Since $\hat{\rho}_{\gamma}(\alpha)$ and $\hat{\rho}_{\gamma}(\beta)$ both contain $\{-\infty\}$, we may find times $s < t$ such that

$$\ell(\beta|\gamma(s)) < e^{-2D} \quad \text{and} \quad \ell(\alpha|\gamma(t)) < e^{-2D}.$$

Corollary 3.7 of [\[BF2\]](#) shows that any primitive loop c with $\ell(c|\gamma(s)) < 1$ (such as β itself) satisfies $\text{diam}_{\mathcal{F}}(\{c\} \cup \pi_{\mathcal{F}}(\gamma(s))) \leq 13$ and consequently $d_{\mathcal{F}}(c, \beta) \leq 30$ by the triangle inequality (recall that $\text{diam}_{\mathcal{F}}(\pi_{\mathcal{F}}(\gamma(s))) \leq 4$ by [\[BF2, Lemma 3.1\]](#)). Evidently, then, we must have $\ell(\alpha|\gamma(s)) \geq 1$. Since γ is a directed geodesic, we thus find that

$$\ell(\alpha|\gamma(u)) \geq e^{-D} > \ell(\alpha|\gamma(t)) \quad \text{for all} \quad u \in [s - D, s].$$

Now, take $0 < \sigma < 1$ sufficiently small (as we may) so that $\pi_{\gamma}(R_{\sigma})$ is contained entirely within $\gamma((-\infty, s - D))$ and so that α is the candidate of R_{σ} that realizes the distance from R_{σ} to $\gamma(t)$. If $[R_{\sigma}, \gamma(t)]$ denotes a directed geodesic from R_{σ} to $\gamma(t)$, it follows that α also realizes the distance from each point $H \in [R_{\sigma}, \gamma(t)]$ to $\gamma(t)$. Thus for every $u \in [s - D, s]$ we have

$$d_{\mathcal{X}}(H, \gamma(t)) = \log \left(\frac{\ell(\alpha|\gamma(t))}{\ell(\alpha|H)} \right) < \log \left(\frac{\ell(\alpha|\gamma(u))}{\ell(\alpha|H)} \right) \leq d_{\mathcal{X}}(H, \gamma(u));$$

consequently $\gamma(u)$ cannot be in $\pi_{\gamma}(H)$. That is, $\pi_{\gamma}([R_{\sigma}, \gamma(t)])$ is disjoint from $\gamma([s - D, s])$. By [Lemma 3.3](#) and the choice of σ , we may conclude that $\pi_{\gamma}([R_{\sigma}, \gamma(t)])$ is contained entirely within $\gamma((-\infty, s - D))$. But this contradicts the obvious fact that $\pi_{\gamma}(\gamma(t)) = \gamma(t)$. The lemma follows. \square

With these basic properties of ρ_{γ} established, we now turn to the main ingredient in the proof of [Theorem 1.3](#). Restricting to the primitive conjugacy classes, our construction $\hat{\rho}_{\gamma}$ (or alternately ρ_{γ}) thus gives a projection $\hat{\rho}_{\gamma}: \mathcal{P}\mathcal{L} \rightarrow \mathcal{P}(\mathbf{I})$ for each geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$. Our next lemma shows that that ρ_{γ} is in fact uniformly Lipschitz provided γ is strongly contracting. Note that this *does not* yet show that the projection ρ_{γ} is a retraction.

Lemma 4.3 (ρ_{γ} is coarsely Lipschitz). *Suppose that $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a D -strongly contracting geodesic and let $\alpha, \beta \in \mathcal{P}\mathcal{L}^0$ be primitive loops. Then $\rho_{\gamma}(\alpha)$ is nonempty (so $\hat{\rho}_{\gamma}(\alpha) \subset \mathbb{R}$) and*

$$\text{diam}_{\mathcal{X}}(\rho_{\gamma}(\alpha) \cup \rho_{\gamma}(\beta)) \leq D \cdot d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) + D.$$

Proof. Let us first prove the following:

Claim. *If $\rho_{\gamma}(\alpha) \neq \emptyset$ and $\beta \in \mathcal{P}\mathcal{L}^0$ is adjacent to α , then $\rho_{\gamma}(\beta) \neq \emptyset$ and $\text{diam}_{\mathcal{X}}(\rho_{\gamma}(\alpha) \cup \rho_{\gamma}(\beta)) \leq D$.*

To prove the claim, we may choose a free basis $\{e_1, \dots, e_n\}$ of \mathbb{F} in which e_1 represents the conjugacy class α and e_2 represents the conjugacy class β . Let (R, g) be the marked rose with petals labeled by the basis elements e_1, \dots, e_n . By [Proposition 2.1](#) there is a finite set \mathcal{C} of candidate conjugacy classes represented by immersed loops in R such that for any metric ℓ on R the distance to any other point $H \in \mathcal{X}$ is given by

$$d_{\mathcal{X}}((R, g, \ell), H) = \log \left(\sup_{z \in \mathcal{C}} \frac{\ell(z|H)}{\ell(z|(R, g, \ell))} \right).$$

Furthermore, both α and β are candidates since they label petals of R .

Choose an arbitrary point $G = \gamma(t) \in \rho_{\gamma}(\alpha)$. If $\rho_{\gamma}(\beta)$ is empty, then $\hat{\rho}_{\gamma}(\beta) \subset \{-\infty, +\infty\}$ meaning that the infimal length m_{β} of β is *only* achieved by sequences of times tending to $\pm\infty$; in which case we may choose $\varepsilon_0 > 0$ sufficiently small so that for any $s_0 \in \mathbf{I}$ we have the implication

$$\ell(\beta|\gamma(s_0)) < m_{\beta} + \varepsilon_0 \implies |s_0 - t| > 2D. \quad (1)$$

We now fix a time parameter $s \in \mathbf{I}$ as follows: If $\rho_{\gamma}(\beta) \neq \emptyset$, then we choose $s \in \hat{\rho}_{\gamma}(\beta)$ arbitrarily; if $\rho_{\gamma}(\beta) = \emptyset$, we instead let $s \in \mathbf{I}$ be any time for which $\ell(\beta|\gamma(s)) < m_{\beta} + \varepsilon_0$. In either case we set $H = \gamma(s)$.

Let us now define

$$K := \max \{ \ell(z|Y) \mid z \in \mathcal{C} \text{ and } Y \in \{G, H\} \} \quad \text{and} \\ k := \min \{ \ell(z|Y) \mid z \in \mathcal{C} \text{ and } Y \in \{G, H\} \}$$

to be the maximal and minimal lengths achieved by any candidate $z \in \mathcal{C}$ at either G or H . Notice that $K \geq k > 0$. Choose a small parameter $0 < \delta < \frac{k}{2rK} \leq \frac{1}{2}$. For each $0 < \sigma < 1$ we let R_{σ} denote the marked metric graph (R, g, ℓ_{σ}) in which the petals of R corresponding to α and β have lengths $\ell(\alpha|R_{\sigma}) = \sigma\delta$ and $\ell(\beta|R_{\sigma}) = (1 - \sigma)\delta$, respectively, and every other petal of R_{σ} has length $\frac{1-\delta}{r-2}$ (so that $\text{vol}(R_{\sigma}) = 1$). Notice that we then have $\ell(z|R_{\sigma}) \geq \frac{1-\delta}{r-2} \geq \frac{1}{2r}$ for *every* candidate $z \in \mathcal{C}$ except for the candidates α and β .

Let us now estimate the distance from R_{σ} to points along $\gamma(\mathbf{I})$. Firstly, at $G \in \gamma(\mathbf{I})$ we have $\ell(\alpha|G) = m_{\alpha}$ and $\ell(z|G) \leq K$ for all other candidates $z \in \mathcal{C}$. Thus we have

$$d_{\mathcal{X}}(R_{\sigma}, G) = \log \sup_{z \in \mathcal{C}} \frac{\ell(z|G)}{\ell(z|R_{\sigma})} \leq \log \sup \left\{ \frac{\ell(\alpha|G)}{\sigma\delta}, \frac{K}{(1-\sigma)\delta}, 2rK \right\}.$$

Since $\ell(\alpha|G) \geq k$ and $\sigma\delta < \delta < \frac{k}{2rK}$, when $\frac{\sigma}{1-\sigma} < \frac{k}{K}$ the above estimate reduces to

$$d_{\mathcal{X}}(R_{\sigma}, G) = \log \left(\frac{\ell(\alpha|G)}{\sigma\delta} \right) = \log \left(\frac{m_{\alpha}}{\sigma\delta} \right).$$

Since m_{α} is the minimal length of α achieved on $\gamma(\mathbf{I})$, we also have

$$d_{\mathcal{X}}(R_{\sigma}, \gamma(u)) \geq \log \left(\frac{\ell(\alpha|\gamma(u))}{\ell(\alpha|R_{\sigma})} \right) \geq \log \left(\frac{m_{\alpha}}{\sigma\delta} \right) = d_{\mathcal{X}}(R_{\sigma}, G) \quad (2)$$

for all $u \in \mathbf{I}$. Therefore $G \in \pi_{\gamma}(R_{\sigma})$ whenever $\frac{\sigma}{1-\sigma} < \frac{k}{K}$.

A similar argument shows that

$$d_{\mathcal{X}}(R_{\sigma}, H) = \log \sup \left\{ \frac{K}{\delta\sigma}, \frac{\ell(\beta|H)}{(1-\sigma)\delta}, 2rK \right\} = \log \left(\frac{\ell(\beta|H)}{(1-\sigma)\delta} \right) \quad (3)$$

whenever $\frac{1-\sigma}{\sigma} < \frac{k}{K}$. If $\rho_\gamma(\beta)$ is nonempty, so that $\ell(\beta|H) = m_\beta$ by choice of H , we again conclude $H \in \pi_\gamma(R_\sigma)$ whenever $\frac{1-\sigma}{\sigma} < \frac{k}{K}$. Otherwise, we note that any closest point $H_\sigma \in \pi_\gamma(R_\sigma)$ satisfies $d_{\mathcal{X}}(R_\sigma, H_\sigma) \leq d_{\mathcal{X}}(R_\sigma, H)$ and consequently $\ell(\beta|H_\sigma) \leq \ell(\beta|H) < m_\beta + \varepsilon_0$ by the choice of H .

Let us now specify parameters $0 < \sigma_\alpha < \sigma_\beta < 1$ by the formulas

$$\frac{\sigma_\alpha}{1-\sigma_\alpha} = \frac{k}{2K} \quad \text{and} \quad \frac{1-\sigma_\beta}{\sigma_\beta} = \frac{k}{2K}.$$

Setting $R_\alpha = R_{\sigma_\alpha}$ and $R_\beta = R_{\sigma_\beta}$, equation (2) ensures that $G \in \pi_\gamma(R_\alpha)$. A comparison of lengths of candidates at R_α and R_β shows that

$$d_{\mathcal{X}}(R_\alpha, R_\beta) = \log \left(\frac{\ell(\alpha|R_\beta)}{\ell(\alpha|R_\alpha)} \right) = \log \left(\frac{\sigma_\beta \delta}{\sigma_\alpha \delta} \right) = \log \left(\frac{2K}{k} \right)$$

and similarly $d_{\mathcal{X}}(R_\beta, R_\alpha) = \log \left(\frac{1-\sigma_\alpha}{1-\sigma_\beta} \right) = \log(2K/k)$. Since this is independent of δ , by choosing δ sufficiently small we can ensure that

$$d_{\mathcal{X}}(R_\alpha, R_\beta) = \log(2K/k) < \log(m_\alpha/\delta\sigma_\alpha) = d_{\mathcal{X}}(R_\alpha, \gamma).$$

Therefore, the D -strongly contracting property implies that

$$\text{diam}_{\mathcal{X}}(\pi_\gamma(R_\alpha) \cup \pi_\gamma(R_\beta)) \leq D. \tag{4}$$

We now finish proving the claim: First consider the case $\rho_\gamma(\beta) \neq \emptyset$, so that $s \in \hat{\rho}_\gamma(\beta)$ by the choice of s and consequently $H = \gamma(s) \in \pi_\gamma(R_\beta)$ by equation (3). Therefore equation (4) ensures $\text{diam}_{\mathcal{X}}(H, G) \leq D$. Since $G = \gamma(t) \in \rho_\gamma(\alpha)$ and $H = \gamma(s) \in \rho_\gamma(\beta)$ were chosen arbitrarily, this proves the claim in the case that $\rho_\gamma(\beta)$ is nonempty. It remains to rule out the possibility $\rho_\gamma(\beta) = \emptyset$, in which case our choice of $H = \gamma(s)$ gives $\ell(\beta|H) < m_\beta + \varepsilon_0$. Let $H' = \gamma(s') \in \pi_\gamma(R_\beta)$ be any closest point from R_β . We then have $d_{\mathcal{X}}(R_\beta, H') \leq d_{\mathcal{X}}(R_\beta, H)$ which, by equation (3), in turn implies

$$\ell(\beta|H') \leq \ell(\beta|H) < m_\beta + \varepsilon_0.$$

Our choice of ε_0 (1) then ensures that $|s' - t| > 2D$. However, since $\gamma(s') \in \pi_\gamma(R_\beta)$ and $\gamma(t) \in \pi_\gamma(R_\alpha)$, this contradicts (4). Thus the contingency $\rho_\gamma(\beta) = \emptyset$ is impossible and the claim holds.

The lemma now easily follows from the claim: Since $\mathcal{P}\mathcal{L}$ is connected and there exists $\alpha_0 \in \mathcal{P}\mathcal{L}^0$ with $\rho_\gamma(\alpha_0) \neq \emptyset$ by Lemma 4.2, the claim shows that $\rho_\gamma(\beta)$ is nonempty for every primitive loop $\beta \in \mathcal{P}\mathcal{L}^0$. Applying the claim inductively with the triangle inequality then gives the desired bound $\text{diam}_{\mathcal{X}}(\rho_\gamma(\alpha) \cup \rho_\gamma(\beta)) \leq D \cdot d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) + D$ for all $\alpha, \beta \in \mathcal{P}\mathcal{L}^0$. \square

5 The progression of thick, strongly contracting geodesics

In this section, we prove our main theorem in the case that the geodesic is contained in some definite thick part of \mathcal{X} . The arguments in this case are made easier by the fact that we can first prove that the diameter of times for which a fixed conjugacy class has bounded length is uniformly controlled. This is the content of Lemma 5.2.

Proposition 5.1 (Thick strongly contracting geodesics make progress in \mathcal{F}). *For each $D \geq 0$ and $\varepsilon > 0$ there exists a constant $K = K(D, \varepsilon) \geq 1$ with the following property. If $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a D -strongly contracting geodesic and $\gamma(\mathbf{I}) \subset \mathcal{X}_\varepsilon$, then $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a K -quasi-geodesic.*

Before proving [Proposition 5.1](#), recall that given a directed geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ and a nontrivial conjugacy class $\alpha \in \mathbb{F}$, we write $m_\alpha = \inf_{t \in \mathbf{I}} \ell(\alpha|\gamma(t))$ for the infimal length that the conjugacy class attains along γ . In the case of a thick strongly contracting geodesic, the set of times where α is short is controlled as follows:

Lemma 5.2 (Transient shortness). *Suppose that $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a D -strongly contracting geodesic with $\gamma(\mathbf{I}) \subset \mathcal{X}_\varepsilon$, and set $\varepsilon' = \varepsilon/(1+2\varepsilon^{-1})$. Then for every primitive element $\alpha \in \mathbb{F}$ we have*

$$\text{diam}_{\mathcal{X}} \left\{ \gamma(s) : s \in \mathbf{I} \text{ and } \ell(\alpha|\gamma(s)) \leq m_\alpha + 2 \right\} \leq 2M_\varepsilon(1 + M_{\varepsilon'}) \log \left(1 + \frac{2}{\varepsilon} \right) + 2D$$

where M_ε is the symmetrization constant provided by [Lemma 2.2](#).

Proof. Suppose that $G, H \in \gamma(\mathbf{I})$ are points for which $\ell(\alpha|G), \ell(\alpha|H) \leq m_\alpha + 2$. Fix a free basis $A = \{e_1, \dots, e_r\}$ of \mathbb{F} with $e_1 = \alpha$ and let (R, g) be the marked rose with petals labeled by elements of A . Let $\mathcal{C} = \mathcal{C}_R$ denote the finite set of candidates of R . For each $0 < \sigma < 1/2$, let $R_\sigma \in \mathcal{X}$ denote the marked metric graph (R, g, ℓ_σ) in which the petal labelled α has $\ell(\alpha|R_\sigma) = \sigma$ and every other petal has length $(1 - \sigma)/(r - 1)$. Notice that \mathcal{C} is the set of candidates for each metric graph R_σ , and that we moreover have $\ell(z|R_\sigma) \geq (1 - \sigma)/(r - 1) \geq \frac{1}{2r}$ for every candidate $z \in \mathcal{C}$ except for α .

We henceforth suppose our parameter satisfies $\sigma < m_\alpha$. By definition of m_α we thus have $\ell(\alpha|\gamma(t))/\ell(\alpha|R_\sigma) \geq \frac{m_\alpha}{\sigma}$ for all $t \in \mathbf{I}$; hence $d_{\mathcal{X}}(R_\sigma, \gamma) \geq \log(m_\alpha/\sigma)$. Now consider the maximum length

$$M = \max_{z \in \mathcal{C}} \{ \ell(z|G), \ell(z|H) \}$$

achieved by any candidate at the two points G, H . If σ is additionally chosen so that $\sigma < m_\alpha/(2rM)$, then we see that for all candidates $\alpha \neq z \in \mathcal{C}$ we have

$$\frac{\ell(\alpha|G)}{\ell(\alpha|R_\sigma)} \geq \frac{m_\alpha}{\sigma} \geq \frac{M}{1/2r} \geq \frac{\ell(z|G)}{\ell(z|R_\sigma)} \quad \text{and} \quad \frac{\ell(\alpha|H)}{\ell(\alpha|R_\sigma)} \geq \frac{m_\alpha}{\sigma} \geq \frac{M}{1/2r} \geq \frac{\ell(z|H)}{\ell(z|R_\sigma)}.$$

It now follows from [Proposition 2.1](#) that

$$\log \left(\frac{m_\alpha}{\sigma} \right) \leq d_{\mathcal{X}}(R_\sigma, \gamma) \leq \left\{ \begin{array}{l} d_{\mathcal{X}}(R_\sigma, G) = \log \frac{\ell(\alpha|G)}{\ell(\alpha|R_\sigma)} \\ d_{\mathcal{X}}(R_\sigma, H) = \log \frac{\ell(\alpha|H)}{\ell(\alpha|R_\sigma)} \end{array} \right\} \leq \log \left(\frac{m_\alpha + 2}{\sigma} \right).$$

Choose a directed geodesic $\rho: [0, K] \rightarrow \mathcal{X}$ from R_σ to G , where $K = d_{\mathcal{X}}(R_\sigma, G)$. Since $e^{d_{\mathcal{X}}(R_\sigma, G)} = \frac{\ell(\alpha|G)}{\ell(\alpha|R_\sigma)}$ and ρ is a geodesic for the Lipschitz metric, it follows that $\ell(\alpha|\rho(t)) = \sigma e^t$ for all $t \in [0, K]$. Hence if we define $G' = \rho(\log(m_\alpha/\sigma))$ (so that $\ell(\alpha|G') = m_\alpha$), we see that

$$d_{\mathcal{X}}(R_\sigma, G') = \log \left(\frac{m_\alpha}{\sigma} \right) \leq d_{\mathcal{X}}(R_\sigma, \gamma) \quad \text{and} \quad d_{\mathcal{X}}(G', G) = \log \left(\frac{\ell(\alpha|G)}{m_\alpha} \right) \leq \log \left(1 + \frac{2}{\varepsilon} \right). \quad (5)$$

Defining H' (on the geodesic from R_σ to H) similarly, we obtain analogous inequalities for H' . By the strongly contracting condition, the first inequality of (5) shows that $\text{diam}_{\mathcal{X}}(\pi_\gamma(G') \cup \pi_\gamma(R_\sigma))$ and similarly $\text{diam}_{\mathcal{X}}(\pi_\gamma(H'), \pi_\gamma(R_\sigma))$ are both bounded by D . Whence

$$\text{diam}_{\mathcal{X}}(\pi_\gamma(G') \cup \pi_\gamma(H')) \leq 2D. \quad (6)$$

On the other hand, the second inequality of (5) shows $G', H' \in \mathcal{X}_{\varepsilon'}$, where $\varepsilon' = \varepsilon/(1+2\varepsilon^{-1})$. Therefore, we also have

$$d_{\mathcal{X}}(G, G'), d_{\mathcal{X}}(H, H') \leq M_{\varepsilon'} \log \left(1 + \frac{2}{\varepsilon} \right).$$

Choose any points $G_0 \in \pi_\gamma(G')$ and $H_0 \in \pi_\gamma(H')$. Since these are by definition *closest* points, $d_{\mathcal{X}}(G', G_0)$ and $d_{\mathcal{X}}(H', H_0)$ can be at most $\log\left(1 + \frac{2}{\varepsilon}\right)$. By the triangle inequality, it follows that

$$d_{\mathcal{X}}(G, G_0), d_{\mathcal{X}}(H, H_0) \leq (1 + M_{\varepsilon'}) \log\left(1 + \frac{2}{\varepsilon}\right).$$

Symmetrizing (Lemma 2.2) to obtain bounds on $\text{diam}_{\mathcal{X}}(G, G_0)$ and $\text{diam}_{\mathcal{X}}(H, H_0)$ and combining with (6), another application of the triangle inequality now gives

$$\text{diam}_{\mathcal{X}}(G, H) \leq 2M_{\varepsilon}(1 + M_{\varepsilon'}) \log\left(1 + \frac{2}{\varepsilon}\right) + 2D. \quad \square$$

Since each primitive loop α in the projection $\pi_{\mathcal{P}\mathcal{L}}(G_t)$ of $G_t = \gamma(t)$ satisfies $\ell(\alpha|G_t) \leq m_{\alpha} + 2$ by definition, Lemma 5.2 shows that the composition

$$\gamma(\mathbf{I}) \xrightarrow{\pi_{\mathcal{P}\mathcal{L}}} \mathcal{P}\mathcal{L} \xrightarrow{\rho_\gamma} \gamma(\mathbf{I})$$

moves points a uniformly bounded distance depending only on D and ε . That is, for each thick strongly contracting geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$, the composition $\pi_{\mathcal{P}\mathcal{L}} \circ \rho_\gamma$ gives a coarse retraction from $\mathcal{P}\mathcal{L}$ onto the image $\pi_{\mathcal{P}\mathcal{L}}(\gamma(\mathbf{I}))$ of γ . Combining this with the fact that ρ_γ is coarsely Lipschitz (Lemma 4.3) now easily implies our main result of this section:

Proof of Proposition 5.1. We write $G_t = \gamma(t)$ for $t \in \mathbf{I}$, and fix $s, t \in \mathbf{I}$ with $s \leq t$. Since the projection $\pi_{\mathcal{F}}: \mathcal{X} \rightarrow \mathcal{F}$ is coarsely 80-Lipschitz [DT, Lemma 2.9] and γ is a geodesic, we immediately have $d_{\mathcal{F}}(G_s, G_t) \leq 80|s - t| + 80$ for all $s, t \in \mathbf{I}$. Thus it remains to bound $d_{\mathcal{F}}(G_s, G_t)$ from below. Let $\alpha \in \mathcal{P}\mathcal{L}^0$ be any primitive conjugacy class represented by an embedded loop in G_s (i.e., any class for which $\alpha|G_s \rightarrow G_s$ is an embedding). Then $\alpha \in \pi_{\mathcal{F}}(G_s)$ by definition of the projection $\pi_{\mathcal{F}}: \mathcal{X} \rightarrow \mathcal{F}$. Similarly choose $\beta \in \mathcal{P}\mathcal{L}^0$ represented by an embedded loop in G_t , so that $\beta \in \pi_{\mathcal{F}}(G_t)$.

Notice that $\ell(\alpha|G_s) \leq 1$ and $\ell(\beta|G_t) \leq 1$ (since the loops are embedded). Thus Lemma 5.2 gives a constant D_{ε} , depending only on ε and D , such that

$$\left. \begin{array}{l} \text{diam}_{\mathcal{X}}(\{G_s\} \cup \rho_\gamma(\alpha)) \\ \text{diam}_{\mathcal{X}}(\{G_t\} \cup \rho_\gamma(\beta)) \end{array} \right\} \leq D_{\varepsilon}$$

Then by Lemma 4.3 we have

$$\begin{aligned} |s - t| &= d_{\mathcal{X}}(G_s, G_t) \\ &\leq \text{diam}_{\mathcal{X}}(\{G_s\} \cup \rho_\gamma(\alpha)) + \text{diam}_{\mathcal{X}}(\rho_\gamma(\alpha) \cup \rho_\gamma(\beta)) + \text{diam}_{\mathcal{X}}(\rho_\gamma(\beta) \cup \{G_t\}) \\ &\leq 2D_{\varepsilon} + D \cdot d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) + D \leq 2D_{\varepsilon} + 2D \cdot d_{\mathcal{F}}(\alpha, \beta) + D \\ &\leq 2D \cdot d_{\mathcal{F}}(G_s, G_t) + 2D_{\varepsilon} + D. \end{aligned}$$

This completes the proof. □

6 Backing into thickness

In light of Proposition 5.1, to prove our main result Theorem 1.3 it now suffices to show that every nondegenerate strongly contracting geodesic γ lives in some definite thick part of \mathcal{X} . We begin by showing that the portion of γ where the lengths of primitive loops are minimized is contained in some definite thick part of \mathcal{X} . Arguments in §§7–8 will then show that all of γ must be thick.

First, recall that L denotes the coarse Lipschitz constant of the projection $\pi_{\mathcal{P}\mathcal{L}}: \mathcal{X} \rightarrow \mathcal{P}\mathcal{L}$. In particular, $d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) \leq L$ for any $\alpha, \beta \in \mathcal{P}\mathcal{L}$ with $\ell(\alpha|G), \ell(\beta|G) \leq 2$ for some $G \in \mathcal{X}$.

Proposition 6.1. *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a D -strongly contracting geodesic and suppose there exist $\alpha_0 \in \mathcal{P}\mathcal{L}$ and $s_0 \in \mathbf{I}$ such that $s_0 \in \hat{\rho}_\gamma(\alpha_0)$ and $\ell(\alpha_0|\gamma(s_0)) \leq 2$. If $\text{diam}_{\mathcal{X}}(\gamma(t_0), \gamma(s_0)) \geq 8DL$ for some $t_0 < s_0$, then*

$$\gamma(\mathbf{I} \cap (-\infty, s_0]) \subset \mathcal{X}_{\varepsilon_0}$$

for some thickness constant $\varepsilon_0 > 0$ depending only on D .

Proof. Suppose that we are given $s_m \in \mathbf{I}$ and $\alpha_m \in \mathcal{P}\mathcal{L}^0$ such that $s_m \in \hat{\rho}_\gamma(\alpha_m)$ and $\ell(\alpha_m|\gamma(s_m)) \leq 2$. Suppose additionally there exists $t_m < s_m$ with $\text{diam}_{\mathcal{X}}(\gamma(t_m), \gamma(s_m)) \geq 8DL$ (note that this holds for $m = 0$). We claim there exists an earlier time $s_{m+1} < s_m$ and a conjugacy class $\alpha_{m+1} \in \mathcal{P}\mathcal{L}^0$ again satisfying the conditions $s_{m+1} \in \hat{\rho}_\gamma(\alpha_{m+1})$ and $\ell(\alpha_{m+1}|\gamma(s_{m+1})) \leq 2$ together with the inequalities

$$\begin{aligned} 4DL &\leq \text{diam}_{\mathcal{X}}(\gamma(s_{m+1}), \gamma(s_m)) \leq D(L + 8DL^2 + 1), \quad \text{and} \\ 3L &\leq d_{\mathcal{P}\mathcal{L}}(\alpha_{m+1}, \alpha_m) \leq L + 8DL^2. \end{aligned} \tag{7}$$

Indeed, by continuity of γ there exists $s'_{m+1} < s_m$ with

$$\text{diam}_{\mathcal{X}}(x'_{m+1}, x_m) = 8DL,$$

where $x_m = \gamma(s_m)$ and $x'_{m+1} = \gamma(s'_{m+1})$. Since γ is a directed geodesic, there exists a candidate α_{m+1} on x'_{m+1} such that $\ell(\alpha_{m+1}|\gamma(s'_{m+1} + t)) = e^t \ell(\alpha_{m+1}|x'_{m+1})$ for all $t \geq 0$. Therefore, if we choose any time $s_{m+1} \in \hat{\rho}_\gamma(\alpha_{m+1})$ realizing the minimal length $m_\gamma(\alpha_{m+1})$, we may be assured that s_{m+1} occurs to the left of s'_{m+1} . Letting $x_{m+1} = \gamma(s_{m+1})$ we thus find that $s_{m+1} \leq s'_{m+1} < s_m$ and $\ell(\alpha_{m+1}|x_{m+1}) \leq \ell(\alpha_{m+1}|x'_{m+1}) \leq 2$, as desired.

To prove the claim it remains to verify the inequalities in (7). First note that $\text{diam}_{\mathcal{X}}(x_{m+1}, x_m) \geq \frac{1}{2} \text{diam}_{\mathcal{X}}(x'_{m+1}, x_m) \geq 4DL$ by the triangle inequality and the fact that γ is a directed geodesic. Hence, using [Lemma 4.3](#) we see that

$$d_{\mathcal{P}\mathcal{L}}(\alpha_{m+1}, \alpha_m) \geq \frac{1}{D} (\text{diam}_{\mathcal{X}}(x_{m+1}, x_m) - D) \geq 3L.$$

On the other hand, we may use the fact that $\ell(\alpha_m|x_m), \ell(\alpha_{m+1}|x'_{m+1}) \leq 2$ to conclude that

$$d_{\mathcal{P}\mathcal{L}}(\alpha_{m+1}, \alpha_m) \leq d_{\mathcal{P}\mathcal{L}}(x'_{m+1}, x_m) \leq L d_{\mathcal{X}}(x'_{m+1}, x_m) + L \leq 8DL^2 + L.$$

Another application of [Lemma 4.3](#) then yields,

$$\begin{aligned} \text{diam}_{\mathcal{X}}(x_{m+1}, x_m) &\leq D \cdot d_{\mathcal{P}\mathcal{L}}(\alpha_{m+1}, \alpha_m) + D \\ &\leq D(L + 8DL^2) + D, \end{aligned}$$

which completes the proof of the claim.

Now let $E := D(L + 8DL^2) + D$ and set

$$\varepsilon_1 = e^{-E} \quad \text{and} \quad \varepsilon_0 = e^{-2E}.$$

We claim that $\gamma([s_{m+1}, s_m]) \subset \mathcal{X}_{\varepsilon_0}$. First observe that if $x_m \notin \mathcal{X}_{\varepsilon_1}$, then we may find $\beta \in \mathcal{P}\mathcal{L}^0$ with $\ell(\beta|x_m) < \varepsilon_1$. In this case (7) would give $\ell(\beta|x_{m+1}) < 1$ showing that β is contained in both projections $\pi_{\mathcal{P}\mathcal{L}}(x_m)$ and $\pi_{\mathcal{P}\mathcal{L}}(x_{m+1})$. However, by (7), this contradicts the fact that these diameter L sets contain α_m and α_{m+1} , respectively. Whence $x_m \in \mathcal{X}_{\varepsilon_1}$ and similarly $x_{m+1} \in \mathcal{X}_{\varepsilon_1}$. Another application of (7) then shows

$$\ell(\beta|\gamma(t)) \geq \ell(\beta|x_m) e^{-|s_m-t|} \geq \varepsilon_1 e^{-E} \geq \varepsilon_0$$

for all $t \in [s_{m+1}, s_m]$. Thus $\gamma([s_{m+1}, s_m]) \subset \mathcal{X}_{\varepsilon_0}$ as claimed.

Let us now prove the proposition. If $\mathbf{I}_- = -\infty$, the above shows that we may find an infinite sequence of times $s_0 > s_1 > \dots$ tending to $-\infty$ such that $\gamma([s_{i+1}, s_i]) \subset \mathcal{X}_{\varepsilon_0}$ for each $i > 0$. Thus the proposition holds in this case. Otherwise $\mathbf{I}_- \neq -\infty$ and we may recursively construct a sequence $s_0 > \dots > s_m$ terminating at a time $s_m \in \mathbf{I}$ for which $\text{diam}_{\mathcal{X}}(\gamma(\mathbf{I}_-), \gamma(s_m)) < 8DL$ and $\gamma(s_m) \in \mathcal{X}_{\varepsilon_1}$ (since $m \geq 1$ by the hypotheses of the proposition). But this implies $\ell(\beta|\gamma(t)) \geq \varepsilon_1 e^{-8DL}$ for every conjugacy class β and time $t \in [\mathbf{I}_-, s_m]$. Thus $\gamma([\mathbf{I}_-, s_m]) \subset \mathcal{X}_{\varepsilon_0}$ as well and the proposition holds. \square

7 Nondegeneracy and thickness

We have now developed enough tools to both establish nondegeneracy for typical strongly contracting geodesics and to show that each nondegenerate strongly contracting geodesic has a uniformly thick initial segment. We first give the proof of [Lemma 3.4](#), showing that strongly contracting geodesics are automatically nondegenerate except possibly in the case of a short geodesic with a very thin left endpoint:

Proof of Lemma 3.4. Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a D -strongly contracting geodesic. First suppose that γ is not infinite to the left (i.e., that $\mathbf{I}_- \neq -\infty$) and let ε be the injectivity radius of $\gamma(\mathbf{I}_-)$. Take $A = M_{\varepsilon'} 18DL$, where $\varepsilon' = \varepsilon e^{-18DL}$. We claim that γ is nondegenerate provided $|\mathbf{I}| \geq A$; this will establish the first item of the lemma. Indeed, consider the points $H = \gamma(\mathbf{I}_-)$ and $G = \gamma(\mathbf{I}_- + A)$. If $G \notin \mathcal{X}_{\varepsilon'}$, then we automatically have $d_{\mathcal{X}}(G, H) \geq \log(\varepsilon/\varepsilon') \geq 18DL$ by definition of the Lipschitz metric, and otherwise $G \in \mathcal{X}_{\varepsilon'}$ so that $d_{\mathcal{X}}(G, H) \geq d_{\mathcal{X}}(H, G)/M_{\varepsilon'} = 18DL$ by [Lemma 2.2](#).

To prove the second item of the lemma, it remains to consider the case $\mathbf{I}_- = -\infty$. Choose $s_0 \in \mathbf{I}$ arbitrarily and let $\alpha \in \mathcal{P}\mathcal{L}^0$ be a primitive conjugacy class with $\ell(\alpha|\gamma(s_0)) \leq 2$ (e.g., a candidate). Next choose a time $s \in \hat{\rho}_{\gamma}(\alpha)$ and note that $\ell(\alpha|\gamma(s)) \leq 2$. The fact that $\mathbf{I}_- = -\infty$ ensures we may find $t < s$ such that $\text{diam}_{\mathcal{X}}(H, G) \geq M_{\varepsilon_0} 18DL$, where $H = \gamma(t)$, $G = \gamma(s)$ and $\varepsilon_0 > 0$ is the thickness constant from [Proposition 6.1](#). Then $\gamma(\mathbf{I} \cap (-\infty, s])$ lies in $\mathcal{X}_{\varepsilon_0}$ by [Proposition 6.1](#), and so we may conclude $d_{\mathcal{X}}(G, H) \geq \text{diam}_{\mathcal{X}}(H, G)/M_{\varepsilon_0} \geq 18DL$ by [Lemma 2.2](#). \square

Our next task is to show that nondegeneracy implies that the hypotheses of [Proposition 6.1](#) are satisfied, and consequently that the initial portion of any such geodesic is uniformly thick. The following lemma will aid in this endeavor.

Lemma 7.1. *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a D -strongly contracting geodesic in \mathcal{X} and suppose that there are $\alpha \in \mathcal{P}\mathcal{L}^0$ and $s, t_1 \in \mathbf{I}$ such that $s \leq t_1$ and $\ell(\alpha|\gamma(t_1)) < e^{-D}\ell(\alpha|\gamma(s))$. Then α has its length minimized to the right of $s \in \mathbf{I}$, i.e. $s < r$ for all $r \in \hat{\rho}_{\gamma}(\alpha)$.*

Proof. Set $H = \gamma(t_1)$ and $J = \mathbf{I} \cap [s - D, s]$. Then α can stretch by at most e^D along J (since γ is a directed geodesic), and so for each $j \in J$ we have

$$\ell(\alpha|\gamma(j)) \geq e^{-D}\ell(\alpha|\gamma(s)) > \ell(\alpha|H).$$

Let $r \in \hat{\rho}_{\gamma}(\alpha)$ be any time minimizing the length of α . Fix a marked rose R with a petal corresponding to the conjugacy class α . For $0 < \sigma < 1$, let R_{σ} denote the metric graph obtained from R by setting the length of the α -petal to σ and the length of each other petal to $\frac{1-\sigma}{r-1}$. As in the proof of [Lemma 4.3](#), σ can be taken sufficiently small so that α is the candidate of R_{σ} realizing the distance from R_{σ} to H . Consequently, if $[R_{\sigma}, H]$ denotes a directed geodesic from R_{σ} to H , then α

also realizes the distance from G to H for each point $G \in [R_\sigma, H]$. It now follows that for each $j \in J$ and $G \in [R_\sigma, H]$ we have

$$d_{\mathcal{X}}(G, \gamma(j)) \geq \log \left(\frac{\ell(\alpha|\gamma(j))}{\ell(\alpha|G)} \right) > \log \left(\frac{\ell(\alpha|H)}{\ell(\alpha|G)} \right) = d_{\mathcal{X}}(G, H).$$

In particular the entire projection $\pi_\gamma([R_\sigma, H])$ is disjoint from the interval $\gamma(J)$.

Taking σ smaller if necessary, we may also assume that α realizes the distance from R_σ to $\gamma(r)$. Since $\ell(\alpha|\gamma(r)) = m_\alpha$ is the minimal length of α , this forces $\gamma(r) \in \pi_\gamma(R_\sigma)$. Whence r cannot lie in J by the above. Now, if $J = \mathbf{I} \cap [s - D, s]$ contains the initial endpoint \mathbf{I}_- , this observation forces $r > s$ as desired. Otherwise J is the length- D interval $J = [s - D, s]$, and we may apply [Lemma 3.3](#) to conclude that $r \in \pi_\gamma([R_\sigma, H])$ is contained in $\gamma(\mathbf{I} \cap (s, \infty))$. Thus $r > s$ and the lemma holds. \square

Corollary 7.2. *Given D there exists $\varepsilon_0 > 0$ with the following property. If $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ a nondegenerate D -strongly contracting geodesic, then there exists $s_0 \in \mathbf{I}$ such that $\gamma(\mathbf{I} \cap (-\infty, s_0]) \subset \mathcal{X}_{\varepsilon_0}$.*

Proof. By definition, nondegeneracy implies that there are times $t_0 < t_1$ in \mathbf{I} so that $d_{\mathcal{X}}(\gamma(t_1), \gamma(t_0)) \geq 18DL$. Letting $s \in \mathbf{I}$ be such that $t_0 \leq s \leq t_1$ and $d_{\mathcal{X}}(\gamma(t_1), \gamma(s)) = 2D$, the triangle inequality then gives

$$d_{\mathcal{X}}(\gamma(s), \gamma(t_0)) \geq d_{\mathcal{X}}(\gamma(t_1), \gamma(t_0)) - 2D \geq 16DL.$$

Let $\alpha \in \mathcal{P}\mathcal{L}$ denote the candidate of $\gamma(t_1)$ that realizes the distance to $\gamma(s)$, i.e. $\ell(\alpha|\gamma(t_1)) = e^{-2D}\ell(\gamma(s))$. If we choose any time $s_0 \in \hat{\rho}_\gamma(\alpha)$ minimizing $\ell(\alpha|\gamma(\cdot))$, then $s \leq s_0$ by [Lemma 7.1](#). Since $\ell(\alpha|\gamma(t_1)) \leq 2$, we have that $\ell(\alpha|\gamma(s_0)) \leq 2$. Finally, since $t_0 < s \leq s_0$ and γ is a directed geodesic, we find that

$$\begin{aligned} 16DL &\leq d_{\mathcal{X}}(\gamma(s), \gamma(t_0)) \leq d_{\mathcal{X}}(\gamma(s), \gamma(s_0)) + d_{\mathcal{X}}(\gamma(s_0), \gamma(t_0)) \\ &\leq d_{\mathcal{X}}(\gamma(t_0), \gamma(s_0)) + d_{\mathcal{X}}(\gamma(s_0), \gamma(t_0)) \leq 2\text{diam}_{\mathcal{X}}(\gamma(t_0), \gamma(s_0)). \end{aligned}$$

Therefore $\text{diam}_{\mathcal{X}}(\gamma(t_0), \gamma(s_0)) \geq 8DL$ and we may apply [Proposition 6.1](#) to complete the proof. \square

Finally, we show that if a strongly contracting geodesic in \mathcal{X} has its initial portion contained in some definite thick part of \mathcal{X} , then the entire geodesic remains uniformly thick.

Lemma 7.3. *Suppose that $\varepsilon_0, D > 0$ and that $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a D -strongly contracting geodesic with $\gamma(\mathbf{I} \cap (-\infty, b]) \subset \mathcal{X}_{\varepsilon_0}$ for some $b \in \mathbf{I}$. Then $\gamma(\mathbf{I}) \subset \mathcal{X}_\varepsilon$, for $\varepsilon = \frac{\varepsilon_0}{2}e^{-4DL}$.*

Proof. Write $G_t = \gamma(t)$ for $t \in \mathbf{I}$. Without loss of generality we assume $\varepsilon_0 < 1$. It suffices to prove $m_\alpha \geq \varepsilon$ where α is an arbitrary primitive loop α . Note that $m_\alpha > 0$ and $\rho_\gamma(\alpha) \neq \emptyset$ by [Lemma 4.3](#). If $m_\alpha \geq \varepsilon_0/2$ there we are done. Otherwise we choose $t_\alpha \in \hat{\rho}_\gamma(\alpha)$ and note that $\ell(\alpha|G_{t_\alpha}) < \varepsilon_0/2$. Since $\ell(\alpha|G_t)$ is continuous in t and at least ε_0 for all $t \leq b$, there is some $s < t_\alpha$ so that $\ell(\alpha|G_s) = \varepsilon_0$.

Let β be the candidate of G_s such that $\ell(\beta|G_{s+t}) = e^t\ell(\beta|G_s)$ for all $t > 0$. If $r \in \hat{\rho}_\gamma(\beta)$ is any time minimizing the length of β , we then necessarily have $r \leq s$. Since α and β each have length less than 2 at $G_s \in \mathcal{X}$, it follows that $d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) \leq L$. [Lemma 4.3](#) then implies that

$$\text{diam}_{\mathcal{X}}(G_r, G_{t_\alpha}) \leq D \cdot d_{\mathcal{P}\mathcal{L}}(\alpha, \beta) + D \leq DL + D \leq 2DL.$$

Since γ is a directed geodesic, $d_{\mathcal{X}}(G_r, G_s) \leq d_{\mathcal{X}}(G_r, G_{t_\alpha})$ and so

$$\begin{aligned} d_{\mathcal{X}}(G_{t_\alpha}, G_s) &\leq d_{\mathcal{X}}(G_{t_\alpha}, G_r) + d_{\mathcal{X}}(G_r, G_s) \\ &\leq d_{\mathcal{X}}(G_{t_\alpha}, G_r) + d_{\mathcal{X}}(G_r, G_{t_\alpha}) \leq 4DL. \end{aligned}$$

In particular, $\frac{\ell(\alpha|G_s)}{\ell(\alpha|G_{t_\alpha})} \leq e^{4DL}$, and so we find $m_\alpha \geq \varepsilon_0 e^{-4DL}$ as desired. \square

8 Characterizing strongly contracting geodesics

We now combine the previous results to complete the proof of our main theorem:

Theorem 1.3 (Strongly contracting geodesics make progress in \mathcal{F}). *For each $D \geq 0$ there exists a constant $K \geq 1$ with the following property. If $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is a nondegenerate D -strongly contracting geodesic, then $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a K -quasigeodesic.*

Proof. Suppose $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ is nondegenerate and D -strongly contracting. By nondegeneracy, [Corollary 7.2](#) and [Lemma 7.3](#) together give $\gamma(\mathbf{I}) \subset \mathcal{X}_\varepsilon$ for some $\varepsilon = \varepsilon(D) > 0$. [Proposition 5.1](#) then provides a constant $K = K(D, \varepsilon)$ for which $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a K -quasigeodesic. \square

We next discuss the converse [Theorem 1.2](#) and explain how the Bestvina–Feighn result [[BF2](#), [Corollary 7.3](#)] on folding lines that make definite progress in \mathcal{F} may be promoted to arbitrary geodesics. While this promotion essentially follows from our earlier work [[DT](#)], we have opted to include a proof here for completeness. In this discussion we assume the reader is familiar with folding paths and standard geodesics in \mathcal{X} ; for background on this see [[FM](#), [BF2](#), [DT](#)].

Theorem 1.2 (Progressing geodesics are strongly contracting). *Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be a geodesic whose projection to \mathcal{F} is a K -quasigeodesic. Then there exists $D \geq 0$ depending only on K (and the injectivity radius of the terminal endpoint of γ) such that γ is D -strongly contracting in \mathcal{X} .*

Proof. Let $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ be an arbitrary directed geodesic such that $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a K -quasigeodesic, and let $H, H' \in \mathcal{X}$ be metric graphs satisfying $d_{\mathcal{X}}(H, H') \leq d_{\mathcal{X}}(H, \gamma)$. [Lemma 4.3](#) of [[DT](#)] shows that $\gamma(\mathbf{I}) \subset \mathcal{X}_\varepsilon$ for some $\varepsilon > 0$ depending only on K (and the injectivity radius of $\gamma(\mathbf{I}_+)$ when $\mathbf{I}_+ \neq +\infty$). Using the coarse symmetry of $d_{\mathcal{X}}$ in \mathcal{X}_ε ([Lemma 2.2](#)), one may easily show that $\pi_\gamma(G)$ is never empty. Hence to prove the theorem it suffices to choose $p \in \pi_\gamma(H)$ and $p' \in \pi_\gamma(H')$ arbitrarily and bound $\text{diam}_{\mathcal{X}}(p, p')$ in terms of K and ε .

Choose a finite subinterval $\mathbf{J} = [a, b] \subset \mathbf{I}$ with $p, p' \in \gamma(\mathbf{J})$ and let $\tilde{\gamma} = \gamma|_{\mathbf{J}}$. Notice that $p \in \pi_{\tilde{\gamma}}(H)$ and $p' \in \pi_{\tilde{\gamma}}(H')$. If $\rho: \mathbf{J} \rightarrow \mathcal{X}$ is any standard geodesic from $\tilde{\gamma}(a)$ to $\tilde{\gamma}(b)$, then [Theorem 4.1](#) of [[DT](#)] ensures $\pi_{\mathcal{F}} \circ \rho$ is a $K' = K'(K, \varepsilon)$ -quasigeodesic and that $\rho(\mathbf{J}) \subset \mathcal{X}_{\varepsilon'}$, where $\varepsilon' = \varepsilon'(K, \varepsilon)$. Consequently, [Proposition 7.2](#) of [[BF2](#)] and [Lemma 4.11](#) of [[DT](#)] (see also [[DT](#), [Proposition 2.10](#)] and the following remark) immediately show that ρ is $D' = D'(K, \varepsilon)$ -strongly contracting.

[Theorem 4.1](#) of [[DT](#)] additionally shows that $\tilde{\gamma}(\mathbf{J})$ and $\rho(\mathbf{J})$ have symmetric Hausdorff distance at most $A' = A'(K, \varepsilon)$. Consequently, we claim that there exists $B' = B'(A', D', \varepsilon')$ such that

$$\text{diam}_{\mathcal{X}}(\pi_{\tilde{\gamma}}(G) \cup \pi_\rho(G)) \leq B' \tag{8}$$

for all $G \in \mathcal{X}$. To see this, choose $Y_0 \in \pi_{\tilde{\gamma}}(G)$ arbitrarily and let $Y \in \pi_\rho(Y_0)$ be a closest point projection of Y_0 to ρ . Noting that $d_{\mathcal{X}}(G, Y_0) \leq d_{\mathcal{X}}(G, \rho) + A'$ and $d_{\mathcal{X}}(Y_0, Y) \leq A'$, we see that $d_{\mathcal{X}}(G, Y) \leq d_{\mathcal{X}}(G, \rho) + 2A'$. Thus, as in the proof of [Lemma 5.2](#), we may find $Y' \in \mathcal{X}$ along a directed geodesic from G to Y such that $d_{\mathcal{X}}(G, Y') \leq d_{\mathcal{X}}(G, \rho)$ and $d_{\mathcal{X}}(Y', Y) \leq 2A'$. The strong contraction property for ρ now gives $\text{diam}_{\mathcal{X}}(\pi_\rho(Y') \cup \pi_\rho(G)) \leq D'$, and the fact that Y is ε' -thick and near Y' bounds $\text{diam}_{\mathcal{X}}(\{Y\} \cup \pi_\rho(Y'))$ in terms of ε' and A' . Hence $\text{diam}_{\mathcal{X}}(\{Y\} \cup \pi_\rho(G))$ is bounded and, since $\text{diam}_{\mathcal{X}}(Y, Y_0) \leq A'$, the claimed inequality (8) holds.

We next claim that π_ρ is coarsely 1-Lipschitz. That is, there exists $C' = C'(D', \varepsilon')$ such that

$$\text{diam}_{\mathcal{X}}(\pi_\rho(G_1) \cup \pi_\rho(G_2)) \leq \text{diam}_{\mathcal{X}}(G_1, G_2) + C' \tag{9}$$

for all $G_1, G_2 \in \mathcal{X}$. Indeed, first consider the case that there exists a directed geodesic $[G_1, G_2]$ with $d_{\mathcal{X}}(Y, \rho) \geq D'$ for all $Y \in [G_1, G_2]$. Dividing $[G_1, G_2]$ into $n = \lceil d_{\mathcal{X}}(G_1, G_2)/D' \rceil$ subgeodesics of

equal length (at most D') and applying strong contraction to each, one finds that

$$\text{diam}_{\mathcal{X}}(\pi_{\rho}(G_1) \cup \pi_{\rho}(G_2)) \leq nD' \leq \left(\frac{d_{\mathcal{X}}(G_1, G_2)}{D'} + 1 \right) D' \leq \text{diam}_{\mathcal{X}}(G_1, G_2) + D'.$$

Next consider the case that $d_{\mathcal{X}}(G_i, \rho) \leq D'$ for each $i = 1, 2$. Choosing $G'_i \in \pi_{\rho}(G_i)$ arbitrarily, [Lemma 2.2](#) and the thickness of $G'_i \in \mathcal{X}_{\varepsilon'}$ together bound $\text{diam}_{\mathcal{X}}(G_i, G'_i)$ in terms of ε' and D' . Thus the difference $|\text{diam}_{\mathcal{X}}(G'_1, G'_2) - \text{diam}_{\mathcal{X}}(G_1, G_2)|$ is bounded in terms of ε' and D' . The general case now follows by subdividing an arbitrary directed geodesic $[G_1, G_2]$ into at most three subgeodesics that each fall under the cases considered above.

To complete the proof of the theorem, note that since $d_{\mathcal{X}}(H, \rho) \geq d_{\mathcal{X}}(H, \bar{\gamma}) - A'$ we can find a point $H_0 \in \mathcal{X}$ (say on a geodesic from H to H') such that $d_{\mathcal{X}}(H, H_0) \leq d_{\mathcal{X}}(H, \rho)$ and $d_{\mathcal{X}}(H_0, H') \leq A'$. Then $\text{diam}_{\mathcal{X}}(\pi_{\rho}(H) \cup \pi_{\rho}(H_0)) \leq D'$ by strong contraction and $\text{diam}_{\mathcal{X}}(\pi_{\rho}(H_0) \cup \pi_{\rho}(H')) \leq A' + C'$ by [\(9\)](#). Combining these with [\(8\)](#) gives the desired bound on $\text{diam}_{\mathcal{X}}(p, p')$. \square

9 Contracting subgroups of $\text{Out}(\mathbb{F})$

In this section we apply [Theorems 1.2–1.3](#) to characterize the finitely generated subgroups of $\text{Out}(\mathbb{F})$ that quasi-isometrically embed into \mathcal{F} . Recall that a subgroup $\Gamma \leq \text{Out}(\mathbb{F})$ is said to be **contracting in \mathcal{X}** if there exists $R \in \mathcal{X}$ and $D > 0$ such that any two points in the orbit $\Gamma \cdot R$ are joined by a D -strongly contracting geodesic (the proof below shows this is in fact equivalent to the following stronger condition: for each $R \in \mathcal{X}$ there exists $D > 0$ such that every directed geodesic between points of $\Gamma \cdot R$ is D -strongly contracting).

Theorem 1.6 (Contracting orbits). *Suppose that $\Gamma \leq \text{Out}(\mathbb{F})$ is finitely generated and that the orbit map $\Gamma \rightarrow \mathcal{X}$ is a quasi-isometric embedding. Then Γ is contracting in \mathcal{X} if and only if the orbit map $\Gamma \rightarrow \mathcal{F}$ to the free factor complex is a quasi-isometric embedding.*

Proof. The “if” direction was essentially obtained by the authors in [\[DT\]](#): Supposing that Γ admits an orbit map into \mathcal{F} that is a quasi-isometric embedding, [Theorem 5.5](#) of [\[DT\]](#) implies that for each $R \in \mathcal{X}$ the orbit $\Gamma \cdot R$ is A -quasiconvex for some $A > 0$. This means that any directed geodesic $\gamma: \mathbf{I} \rightarrow \mathcal{X}$ between orbit points lies in the symmetric A -neighborhood of $\Gamma \cdot R$. Since $\Gamma \rightarrow \mathcal{F}$ is a quasi-isometric embedding, it follows easily that the projection $\pi_{\mathcal{F}} \circ \gamma: \mathbf{I} \rightarrow \mathcal{F}$ is a parameterized quasigeodesic with uniform constants. Therefore γ is uniformly strongly contracting by [Theorem 1.2](#).

For the “only if” direction, suppose that Γ is contracting with respect to $R \in \mathcal{X}$ and $D > 0$ and that the assignment $g \mapsto g \cdot R$ defines a C -quasi-isometric embedding. Choose $g, h \in \Gamma$ and let $\gamma: [a, b] \rightarrow \mathcal{X}$ be a D -strongly contracting geodesic from $g \cdot R$ to $h \cdot R$. [Lemma 7.3](#) then ensures $\gamma([a, b]) \subset \mathcal{X}_{\varepsilon}$ for some $\varepsilon > 0$ depending on D and the injectivity radius of R , and so [Proposition 5.1](#) implies that $\pi_{\mathcal{F}} \circ \gamma$ is a $K = K(D, \varepsilon)$ -quasigeodesic. Since $d_{\Gamma}(g, h)$ and $d_{\mathcal{F}}(g\pi_{\mathcal{F}}(R), h\pi_{\mathcal{F}}(R))$ both coarsely agree with $d_{\mathcal{X}}(\gamma(a), \gamma(b)) = d_{\mathcal{X}}(g \cdot R, h \cdot R)$, there is a constant $E = E(K, C) \geq 1$ such that

$$\frac{1}{E} d_{\Gamma}(g, h) - E \leq \text{diam}_{\mathcal{F}}(g\pi_{\mathcal{F}}(R), h\pi_{\mathcal{F}}(R)) \leq E d_{\Gamma}(g, h) + E$$

Thus the assignment $g \mapsto g \cdot A$, where $A \in \pi_{\mathcal{F}}(R)$, defines a quasi-isometric embedding $\Gamma \rightarrow \mathcal{F}$. \square

References

- [AK] Yael Algom-Kfir. Strongly contracting geodesics in Outer space. *Geom. Topol.*, 15:2181–2233, 2011.

- [AKB] Yael Algom-Kfir and Mladen Bestvina. Asymmetry of outer space. *Geom. Dedicata*, 156(1):81–92, 2012.
- [BF1] Mladen Bestvina and Mark Feighn. A hyperbolic $\text{Out}(F_n)$ -complex. *Groups, Geometry, and Dynamics*, 4(1):31–58, 2010.
- [BF2] Mladen Bestvina and Mark Feighn. Hyperbolicity of the complex of free factors. *Adv. Math.*, 256:104–155, 2014.
- [BR] Mladen Bestvina and Patrick Reynolds. The boundary of the complex of free factors. Preprint arXiv:1211.3608, 2012.
- [CV] Marc Culler and Karen Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.*, 84(1):91–119, 1986.
- [DT] Spencer Dowdall and Samuel J Taylor. Hyperbolic extensions of free groups. Preprint arXiv:1406.2567, 2014.
- [FM] Stefano Francaviglia and Armando Martino. Metric properties of outer space. *Publ. Mat.*, 55(2):433–473, 2011.
- [Ham1] Ursula Hamenstädt. The boundary of the free factor graph and the free splitting graph. Preprint arXiv:1211.1630, 2013.
- [Ham2] Ursula Hamenstädt. Lines of minima in Outer space. *Duke Math. J.*, 163:733–776, 2014.
- [HH] Ursula Hamenstädt and Sebastian Hensel. Convex cocompact subgroups of $\text{Out}(F_n)$. Preprint arXiv:1411.2281, 2014.
- [HM] Michael Handel and Lee Mosher. The expansion factors of an outer automorphism and its inverse. *Trans. Amer. Math. Soc.*, 359(7):3185–3208, 2007.
- [Hor] Camille Horbez. The Poisson boundary of $\text{Out}(F_n)$. Preprint arXiv:1405.7938, 2014.
- [HV] Allen Hatcher and Karen Vogtmann. The complex of free factors of a free group. *Quart. J. Math.*, 49(196):459–468, 1998.
- [Min] Yair N. Minsky. Quasi-projections in Teichmüller space. *J. Reine Angew. Math.*, 473:121–136, 1996.
- [Pau] Frédéric Paulin. The Gromov topology on R-trees. *Topology Appl.*, 32(3):197–221, 1989.

Department of Mathematics
 University of Illinois at Urbana-Champaign
 1409 W. Green Street
 Urbana, IL 61801, U.S.A
 E-mail: dowdall@illinois.edu

Department of Mathematics
 Yale University
 10 Hillhouse Ave
 New Haven, CT 06520, U.S.A
 E-mail: s.taylor@yale.edu